

Finite size effects in a stochastic condensation model


Paul Chleboun
Stefan Grosskinsky

Complexity Science Doctoral Training Centre
University of Warwick

29th October 2009

Model: The zero-range process

A continuous-time Markov Chain. Driven diffusive system.



[Spitzer (1970), Andjel (1982)]

Lattice: $\Lambda_L = \{1, \dots, L\}$ with pbc

Configuration: $\eta = (\eta_x)_{x \in \Lambda_L}$ with $\eta_x \in \mathbb{N}_0$

State space: $X_L = \mathbb{N}_0^{\Lambda_L}$

Jump rates: $g : \mathbb{N}_0 \rightarrow [0, \infty)$

$$g(n) = 0 \iff n = 0$$


Motivation

$g(k) \searrow \Rightarrow$ effective attraction, condensation possible

$$g(k) \simeq 1 + \frac{b}{k^\gamma}, \quad b > 0, \gamma \in (0, 1)$$




[Evans (2000)]

- Many applications. [Evans and Hanney (2005)]
- Including to **granular media** and **traffic flow**.



[van der Meer, van der Weele, Lohse, Mikkelsen, Versluis (2001-02)]

Motivation: Granular media




[van der Meer, van der Weele, Lohse, Mikkelsen, Versluis (2001-02)]
stilton.tnw.utwente.nl/people/rene/clustering.html

- Typical sizes $L \sim 10 - 100$, $N \sim 1000$.

Motivation: Traffic

Mapping **ZRP** to **ASEP**:



- ASEP can be considered as a simple traffic model

[Kaupuzs, Mahnke, Harris (2008)]

- Typical sizes $L \sim 1000$, $N \sim 200$.

Outline

- 1 Introduction
 - Model: The zero-range process
 - Heuristic Motivation
 - Previous Results: Equivalence of ensembles
- 2 Observations
 - Numerics
 - MC simulations
- 3 Results
 - Fluid and Condensed approximations
 - Rate function
 - Current matching
 - Scaling limit
- 4 Conclusions

Stationary measures: Grand canonical ensemble

There exist stationary product measures.

Stationary weights: $w(n) = \prod_{k=1}^n g(k)^{-1}$

Grand canonical ensemble

$$\nu_{\phi}^L(\eta) = \frac{1}{z(\phi)^L} \prod_{x \in \Lambda_L} w(\eta_x) \phi^{\eta_x} \quad \text{where} \quad z(\phi) = \sum_{n=0}^{\infty} w(n) \phi^n$$

Fugacity: $\phi \in [0, \phi_c)$

(↗) **Density:** $\langle \eta_x \rangle_{\nu_{\phi}} = R(\phi) = \phi \partial_{\phi} \log z(\phi), \quad \rho_c = \lim_{\phi \rightarrow \phi_c} R(\phi) \in (0, \infty]$

Current: $\langle g(\eta_x) \rangle_{\nu_{\phi}} = \phi$

Stationary measures: Canonical ensemble

Dynamics conserve the total particle number:

$$\Sigma_L(\boldsymbol{\eta}) := \sum_{x \in \Lambda_L} \eta_x = N$$

Fixed L and N : system is irreducible and finite state so **ergodic**.

Unique stationary measure:

Canonical Ensemble

$$\pi_{L,N}(\boldsymbol{\eta}) := \frac{1}{Z(L,N)} \prod_{x \in \Lambda_L} w(\eta_x) \delta\left(\sum_x \eta_x - N\right)$$

Density: $\langle \eta_x \rangle_{\pi_{L,N}} = \rho = N/L$

Thermodynamic limit

Equivalence of ensembles:




Previous results

[Großkinsky, Schütz, Spohn (2003)]

In the thermodynamic limit $L, N \rightarrow \infty$, $N/L \rightarrow \rho$

$$\pi_{L,N} \xrightarrow{w} \nu_{\Phi(\rho)} \quad \text{where} \quad \Phi(\rho) = \begin{cases} R^{-1}(\rho) & \text{if } \rho < \rho_c \\ \phi_c & \text{if } \rho \geq \rho_c \end{cases}$$

$$g(k) = 1 + \frac{b}{k^\gamma}, \quad b > 0, \gamma \in (0, 1) \implies \rho_c < \infty$$





Thermodynamic limit

Thermodynamic entropy:

$$s(\rho) = \sup_{\phi \in [0,1]} (\log z(\phi) - \rho \log \phi)$$

Convergence of canonical entropy:

$$\lim_{L \rightarrow \infty} \frac{1}{L} \log Z(L, [\rho L]) = \log z(\Phi(\rho)) - \rho \log \Phi(\rho) = s(\rho)$$



$$\Phi(\rho) = e^{-\partial_{\rho} s(\rho)}$$


Numerics in the Canonical Ensemble

Recursion relation

$$Z(L, N) = \sum_{k=0}^N w(k) Z(L-1, N-k)$$

Canonical current



$$j := \langle g(\eta_x) \rangle_{\pi_{L,N}} = \frac{Z(L, N-1)}{Z(L, N)}$$



- Large fluid current overshoot.
- Sharp transition to putative condensed phase.



Numerics in the Canonical Ensemble

Exact numerics, canonical current

 $\gamma = 0.5, b = 4$ 



$$s_L(\rho) = \frac{1}{L} \log Z(L, N)$$

Monte Carlo simulations



- Left: Metastable switching between fluid and condensed currents.
- Right: Exponential distribution of waiting time in the condensed and fluid ‘phases’.

Fluid and Condensed approximations



- Describe where the approximations come from
- How they can be used to find the leading order effect

Rate function

Aim:

- Find the scaling of the overshoot region.
- Understand the apparent metastability close to the maximum current .

Approach:

- Find something like a rate function which describes the metastability in some scaling limit in terms of the background density.

$$\pi_{L,N} \left(\frac{N - \max_{x \in \Lambda} \eta_x}{L - 1} \asymp \rho_{\text{bg}} \right) \sim \exp \left(-L^\alpha I_\rho(\rho_{\text{bg}}) \right)$$

Rate function

Notation:

$$\begin{aligned} \pi_{L,N}(\rho_{\text{bg}}) &:= \pi_{L,N} \left(N - \max_{x \in \Lambda_L} \eta_x = \rho_{\text{bg}}(L-1) \right) \\ &= \pi_{L,N} \left(\max_{x \in \Lambda_L} \eta_x = \eta_{\text{max}} \right) \end{aligned}$$

where $\rho_{\text{bg}}(L-1) = N - \eta_{\text{max}}$.

Rate function

Notation:

$$\begin{aligned}\pi_{L,N}(\rho_{\text{bg}}) &:= \pi_{L,N} \left(N - \max_{x \in \Lambda_L} \eta_x = \rho_{\text{bg}}(L-1) \right) \\ &= \pi_{L,N} \left(\max_{x \in \Lambda_L} \eta_x = \eta_{\text{max}} \right)\end{aligned}$$

where $\rho_{\text{bg}}(L-1) = N - \eta_{\text{max}}$.

We have:

$$\pi_{L,N}(\rho_{\text{bg}}) = \frac{1}{Z(L,N)} L w(\eta_{\text{max}}) \sum_{\eta \in \tilde{X}} \prod_{x=1}^{L-1} w(\eta_x)$$

where $\tilde{X} = \left\{ \eta : \sum_{x=1}^{L-1} \eta_x = \rho_{\text{bg}}(L-1), \eta_1, \dots, \eta_{L-1} \leq N - \rho_{\text{bg}}(L-1) \right\}$.

Rate function

$$\tilde{X} = \left\{ \boldsymbol{\eta} : \sum_{x=1}^{L-1} \eta_x = \rho_{\text{bg}}(L-1), \eta_1, \dots, \eta_{L-1} \leq N - \rho_{\text{bg}}(L-1) \right\}$$

Rate function

$$\tilde{X} = \left\{ \boldsymbol{\eta} : \sum_{x=1}^{L-1} \eta_x = \rho_{\text{bg}}(L-1), \eta_1, \dots, \eta_{L-1} \leq N - \rho_{\text{bg}}(L-1) \right\}$$

$$\begin{aligned} \pi_{L,N}(\rho_{\text{bg}}) &= \frac{1}{Z(L,N)} Lw(\eta_{\text{max}}) \sum_{\boldsymbol{\eta} \in \tilde{X}} \prod_{x=1}^{L-1} w(\eta_x) \\ &= \frac{1}{Z(L,N)} Lw(\eta_{\text{max}}) \phi^{-\rho_{\text{bg}}(L-1)} \sum_{\boldsymbol{\eta} \in \tilde{X}} \prod_{x=1}^{L-1} w(\eta_x) \phi^{\eta_x} \\ &= \frac{1}{Z(L,N)} Lw(\eta_{\text{max}}) \phi^{-\rho_{\text{bg}}(L-1)} \mathcal{Z}_{\eta_{\text{max}}}^{L-1}(\phi) \nu_{\phi, \eta_{\text{max}}}^{L-1}(\Sigma_{L-1} = \rho_{\text{bg}}(L-1)) \end{aligned}$$

Rate function

$$\tilde{X} = \left\{ \boldsymbol{\eta} : \sum_{x=1}^{L-1} \eta_x = \rho_{\text{bg}}(L-1), \eta_1, \dots, \eta_{L-1} \leq N - \rho_{\text{bg}}(L-1) \right\}$$

$$\begin{aligned} \pi_{L,N}(\rho_{\text{bg}}) &= \frac{1}{Z(L,N)} Lw(\eta_{\text{max}}) \sum_{\boldsymbol{\eta} \in \tilde{X}} \prod_{x=1}^{L-1} w(\eta_x) \\ &= \frac{1}{Z(L,N)} Lw(\eta_{\text{max}}) \phi^{-\rho_{\text{bg}}(L-1)} \sum_{\boldsymbol{\eta} \in \tilde{X}} \prod_{x=1}^{L-1} w(\eta_x) \phi^{\eta_x} \\ &= \frac{1}{Z(L,N)} Lw(\eta_{\text{max}}) \phi^{-\rho_{\text{bg}}(L-1)} \mathcal{Z}_{\eta_{\text{max}}}^{L-1}(\phi) \nu_{\phi, \eta_{\text{max}}}^{L-1}(\sum_{L-1} = \rho_{\text{bg}}(L-1)) \end{aligned}$$

where,

$$\nu_{\phi, \eta_{\text{max}}}^{L-1}(\sum_{L-1} = \rho_{\text{bg}}(L-1)) = \nu_{\phi}^{L-1}(\sum_{L-1} = \rho_{\text{bg}}(L-1) | \eta_x \leq \eta_{\text{max}})$$

Rate function

$$\begin{aligned} \log \pi_{N,L}(\rho_{\text{bg}}) &= (L-1) (\log z_{\eta_{\max}}(\phi) - \rho_{\text{bg}} \log \phi) + \\ &\quad + \log w(\eta_{\max}) + \log \nu_{\phi, \eta_{\max}}^{L-1}(\sum_{L-1} = \rho_{\text{bg}}(L-1)) - \\ &\quad - \log Z(L, N) + \log L. \end{aligned}$$

Holds for all $\phi \in [0, \infty)$. Choose ϕ so that,

$$R_{\eta_{\max}}(\phi) := \frac{1}{z_{\eta_{\max}}(\phi)} \sum_{k=1}^{\eta_{\max}} kw(k)\phi^k = \rho_{\text{bg}}$$

Rate function

$$\begin{aligned} \log \pi_{N,L}(\rho_{\text{bg}}) &= (L-1) \left(\log z_{\eta_{\max}}(\Phi_{\eta_{\max}}(\rho_{\text{bg}})) - \rho_{\text{bg}} \log \Phi_{\eta_{\max}}(\rho_{\text{bg}}) \right) + \\ &\quad + \log w(\eta_{\max}) + \log \nu_{\Phi_{\eta_{\max}}(\rho_{\text{bg}}), \eta_{\max}}^{L-1}(\sum_{L-1} = \rho_{\text{bg}}(L-1)) - \\ &\quad - \log Z(L, N) + \log L. \end{aligned}$$

Holds for all $\phi \in [0, \infty)$. Choose ϕ so that,

$$R_{\eta_{\max}}(\phi) := \frac{1}{z_{\eta_{\max}}(\phi)} \sum_{k=1}^{\eta_{\max}} k w(k) \phi^k = \rho_{\text{bg}}$$

Define: $\Phi_{\eta_{\max}} = R_{\eta_{\max}}^{-1}$.

Rate function

$$\begin{aligned} \log \pi_{N,L}(\rho_{\text{bg}}) = & (L-1) \left(\log z_{\eta_{\text{max}}}(\Phi_{\eta_{\text{max}}}(\rho_{\text{bg}})) - \rho_{\text{bg}} \log \Phi_{\eta_{\text{max}}}(\rho_{\text{bg}}) \right) + \\ & + \log w(\eta_{\text{max}}) + \log \nu_{\Phi_{\eta_{\text{max}}}(\rho_{\text{bg}}), \eta_{\text{max}}}^{L-1}(\Sigma_{L-1} = \rho_{\text{bg}}(L-1)) - \\ & - \log Z(L, N) + \log L. \end{aligned}$$

- Looks like the thermodynamic entropy, except with truncation at η_{max} so exists for all ρ_{bg} .

Rate function

$$\begin{aligned} \log \pi_{N,L}(\rho_{\text{bg}}) = & (L-1) \left(\log z_{\eta_{\text{max}}}(\Phi_{\eta_{\text{max}}}(\rho_{\text{bg}})) - \rho_{\text{bg}} \log \Phi_{\eta_{\text{max}}}(\rho_{\text{bg}}) \right) + \\ & + \log w(\eta_{\text{max}}) + \log \nu_{\Phi_{\eta_{\text{max}}}(\rho_{\text{bg}}), \eta_{\text{max}}}^{L-1}(\Sigma_{L-1} = \rho_{\text{bg}}(L-1)) - \\ & - \log Z(L, N) + \log L. \end{aligned}$$

- Looks like the thermodynamic entropy, except with truncation at η_{max} so exists for all ρ_{bg} .
- Contribution due to the condensate.

Rate function

$$\begin{aligned} \log \pi_{N,L}(\rho_{\text{bg}}) = & (L-1) \left(\log z_{\eta_{\text{max}}}(\Phi_{\eta_{\text{max}}}(\rho_{\text{bg}})) - \rho_{\text{bg}} \log \Phi_{\eta_{\text{max}}}(\rho_{\text{bg}}) \right) + \\ & + \log w(\eta_{\text{max}}) + \log \nu_{\Phi_{\eta_{\text{max}}}(\rho_{\text{bg}}), \eta_{\text{max}}}^{L-1}(\Sigma_{L-1} = \rho_{\text{bg}}(L-1)) - \\ & - \log Z(L, N) + \log L. \end{aligned}$$

- Looks like the thermodynamic entropy, except with truncation at η_{max} so exists for all ρ_{bg} .
- Contribution due to the condensate.
- Can be approximated by Gaussian density at 0.


Rate function

$$\begin{aligned} \log \pi_{N,L}(\rho_{\text{bg}}) = & (L-1) \left(\log z_{\eta_{\text{max}}}(\Phi_{\eta_{\text{max}}}(\rho_{\text{bg}})) - \rho_{\text{bg}} \log \Phi_{\eta_{\text{max}}}(\rho_{\text{bg}}) \right) + \\ & + \log w(\eta_{\text{max}}) + \log \nu_{\Phi_{\eta_{\text{max}}}(\rho_{\text{bg}}), \eta_{\text{max}}}^{L-1}(\Sigma_{L-1} = \rho_{\text{bg}}(L-1)) - \\ & - \log Z(L, N) + \log L. \end{aligned}$$

- Looks like the thermodynamic entropy, except with truncation at η_{max} so exists for all ρ_{bg} .
- Contribution due to the condensate.
- Can be approximated by Gaussian density at 0.
- The final terms are constant for fixed N and L .

Rate function


Ignoring the final two normalising terms we can approximate $-\log \pi_{N,L}(\rho_{\text{bg}})$ very quickly. Still informative.



- ρ below sharp transition point.

Rate function


Ignoring the final two normalising terms we can approximate $-\log \pi_{N,L}(\rho_{\text{bg}})$ very quickly. Still informative.



- ρ near sharp transition point.

Rate function

Ignoring the final two normalising terms we can approximate $-\log \pi_{N,L}(\rho_{\text{bg}})$ very quickly. Still informative.



- ρ above sharp transition point.


Max and min of distribution

Under controllable approximations (to leading order in L):


$$\log \pi_{N,L}(\rho_{\text{bg}}) - \log \pi_{N,L}(\rho_{\text{bg}} - 1/L) \simeq \log g(\eta_{\text{max}}) - \log \Phi_{\eta_{\text{max}}}(\rho_{\text{bg}})$$

- First minimum and maximum are given by current matching.
- The final minimum is effectively a boundary minimum.

Current matching




Lifetime of fluid and condensed 'phases'



Fixed Density N/L

- Ratio of the areas will always give proportion of time spent in each 'phase'
- For totally asymmetric process random walk argument very well to predict the life time.

Approximations



- **Fluid current:** From the average current in a cut-off grand canonical ensemble $\Phi_N(\rho)$.
- **Condensed current:** From current matching, solve

$$\Phi_N(\rho_{bg}) = g(\eta_{max})$$

Scaling limit


Current overshoot is asymptotically linear:

$$(R_{n_L}(\phi) - \rho_c) \sim \sigma_c^2(\phi - \phi_c) \quad \text{so,} \quad (\Phi_N(\rho_{\text{bg}}) - 1) \sim \frac{1}{\sigma_c^2}(\rho_{\text{bg}} - \rho_c)$$

where $\sigma_c^2 = \xi_c - \rho_c^2$ and $\xi_c = \sum_{k=0}^{\infty} k^2 \nu_{\phi_c}(k)$


- This leads to the asymptotic behaviour of the current matching curve $\Phi_N(\rho_{\text{bg}}) = g(\eta_{\text{max}})$.
- From this we can derive the scaling so that the current matching curves collapse.

Scaling limit




- Black: Numerics average current, $L = 10000$ system
- Blue: Linear fluid overshoot
- Red: Collapsed current matching curve

Current matching




- Blue: Linear scaling limit for fluid current in background.
- Red: Current out of condensate (Collapsed in scaling limit).

Current matching



- Blue: Linear scaling limit for fluid current in background.
- Red: Current out of condensate (Collapsed in scaling limit).
- These give a unique non-degenerate rate function in the scaling limit.


Scaling rate function



$$I_{\tilde{\delta\rho}}(\delta\tilde{\rho}_{bg}) := \lim_{L \rightarrow \infty} \frac{1}{L^\alpha} \log \pi_{L, [(\rho_c + \delta\rho L^{-\alpha})L]} \left(\rho_{bg} \asymp \rho_c + \delta\tilde{\rho}_{bg} L^{-\alpha} \right)$$


where $\alpha = \frac{\gamma}{\gamma + 1}$.

Transition point



- Transition point from ‘fluid’ to ‘condensed’ is given by the rescaled density for which minimum are equal depth.
- Find using rescaled entropies from integrating the two current curves.
- ‘Fluid’ and ‘condensed’ entropies cross at the transition.

Data collapses



- Also for each vertical slice (fixed rescaled density) we have the scaling rate function.

Conclusions and Summary

- Large finite size effects observed for some parameter values.
- Observed phenomena on a finite system such as metastable switching are also observed in real world clustering.
- May have implications for understanding traffic flow patterns.

Results:

- Estimate the fluid current overshoot and condensed current on large finite systems.
- Estimate the density at which an observed change in 'phase' will occur on a large finite system.
- Predict asymptotic scaling.
- Predict switching times in terms of entropy differences (for certain systems).

Acknowledgements

Thanks to

- Stefan Grosskinsky for supervision.
- Second supervisor Ellak Somfai.
- Discussions with Robin Ball.
- Warwick Complexity Science DTC
- Funding from the EPSRC