

Thames study: Rivers can be a source antibiotic resistance

Feb 13, 2015



Image: USGS

Rivers and streams could be a major source of antibiotic resistance in the environment.

The discovery comes following a study on the Thames river by scientists at the University of Warwick's School of Life Sciences and the University of Exeter Medical School.

The study found that greater numbers of [resistant bacteria](#) exist close to some [waste water](#) treatment works, and that these plants are likely to be responsible for at least half of the increase observed.

Antimicrobial resistance is one of the largest threats to human health for a century, the researchers argue. Increasingly large amounts of antibiotics are released into the environment through both human and agricultural use, with surface run off from farming activities (including fertiliser and animal slurry) washed straight into rivers after [heavy rainfall](#).

Co-lead on the research, Professor Elizabeth Wellington of the University of Warwick, said:

"Antibiotic resistance naturally occurs in the environment, but we don't yet know how human and agricultural waste is affecting its development. We've found that waste water discharges effect resistance levels and that improvements in our treatment processes could hold the key to reducing the prevalence of resistant bacteria in the environment.

"We found antibiotic resistance in the group Enterobacteriaceae which includes gut bacteria and pathogens."

Published in Nature's *The ISME Journal*, the study has also shown that different types of [waste water treatment](#) plant release varying amounts of resistant bacteria. Professor Wellington explains:

We produced a model based on our data which showed that there was a big difference between secondary and tertiary activated sludge plant where the latter resulted in a predicted 100-fold decrease in resistance levels.


Study co-lead author, Dr William Gaze of the University of Exeter Medical School said:

"Our research has shed further light on links between environmental pollutants and antibiotic resistance. It has allowed us to uncover an association between a number of compounds - such as zinc, phosphorous and silicon - and antibiotic resistance. We think those bacteria that have developed to survive in environments rich in metals may also possess antibiotic resistance mechanisms - highlighting the complexity of this

Featured

Popular

Most shared



[Iconic graph at center of climate debate](#) Feb 14, 2015 81



[Possible discovery in 2015 of a new particle in physics](#) 18 hours ago 25



[Where did the Big Bang happen?](#) Feb 13, 2015 53



[Interstellar technology throws light on spinning black holes](#) Feb 13, 2015 76



[Over 100 banks hit by sophisticated cyberattack: report](#) Feb 14, 2015 24

Phys.org on facebook



You and 913,789 others like this. 913,789 people

Relevant PhysicsForums posts

[Earth's inner core has an inner core of its own](#) 17 hours ago

[Visually stunning minerals/crystals](#) Feb 14, 2015

[Enthalpy of condensation and the global heat budget](#) Feb 14, 2015

[Dirty rain in the Northwest](#) Feb 12, 2015

["Why models run hot"](#) Feb 11, 2015

global issue."

The researchers analysed water and sediment samples from 13 sites across the Thames river catchment and developed detailed models to predict the distribution of [antibiotic resistant bacteria](#).

The team also found that several other factors affected the prevalence of antibiotic resistance, such as changes in rainfall and land cover. For example, heavy rainfall at a point surrounded by grassland raised resistance levels; whereas a heavy rainfall at a point surrounded by woodland reduced the levels seen.

The findings have allowed the research team to develop a robust model that will predict the level of antibiotic resistance in other catchments, without the need for detailed water sampling.

Increased levels of [antibiotic resistance](#) in the aquatic environment could lead to increased risk of human exposure. More research is required to fully understand the risk posed via this route and the possible implications for public health.

The study is published in the journal of the *International Society for Microbial Ecology*.

+ Explore further: [Sewage treatment contributes to antibiotic resistance](#)

Journal reference: [ISME Journal](#)

Provided by [University of Warwick](#)



5 / 5 (7 votes)

[Large crater in Siberia](#) Feb 09, 2015

More from [Earth](#)

Related Stories



[Sewage treatment contributes to antibiotic resistance](#) Jul 21, 2014

Wastewater treatment plants are unwittingly helping to spread antibiotic resistance, say scientists.

[Obama to seek more funds for antibiotic-resistant bacteria fight](#) Jan 27, 2015


President Barack Obama will request a doubling of funds for fighting and preventing antibiotic-resistant bacteria in his 2016 budget, the White House said Tuesday.



[Environmental scientists find antibiotics, bacteria, resistance genes in feedlot dust](#) Jan 23, 2015

After testing dust in the air near cattle feedlots in the Southern High Plains, researchers at The Institute of

Recommended for you



[Satellite images reveal ocean acidification from space](#) 3 hours ago

Pioneering techniques that use satellites to monitor ocean acidification are set to revolutionise the way that marine biologists and climate scientists study the ocean. This new approach, that will be published ...



[Scientists seek international authority on climate geoengineering](#) 18 hours ago

US scientists and legal experts are calling for a strong, international authority to regulate any man-made interventions meant to combat global warming, amid fears that the technology could be harmful to ...



[Limited climate change accord likely in Paris, IPCC expert says](#) 18 hours ago

1 comment



Adjust slider to filter visible comments by rank

Display comments: **newest first**

Returners **★** not rated yet

Feb 14, 2015



whereas a heavy rainfall at a point surrounded by woodland reduced the levels seen.