

MA231 Vector Analysis

Example Sheet 2: Hints and partial solutions

2009, term 1
Stefan Adams

A1 (a) $4x$ (b) $y(1 + x^2y^2z^2)^{-1/2}$ (c) xye^z .

A2 (a) $\hat{N} = \frac{x}{R}$ on the surface \mathcal{S} so that $v \cdot \hat{N} = \|x\|^2/R = R$ and $\int_{\mathcal{S}} v \cdot \hat{N} \, dS = R \int_{\mathcal{S}} dS = R$ (Surface area of \mathcal{S}) $= 4\pi R^3$. (b) Here $v \cdot \hat{N} = 0$ so the flux is zero. (c) Here $v \cdot \hat{N} = \frac{1}{R}(-x^2 + y^2 + z^2)$. But by symmetry $\int_{\mathcal{S}} x^2 \, dS = \int_{\mathcal{S}} y^2 \, dS = \int_{\mathcal{S}} z^2 \, dS$. So $\int_{\mathcal{S}} v \cdot \hat{N} \, dS = \frac{1}{R} \int_{\mathcal{S}} z^2 \, dS = \frac{1}{3R} \int_{\mathcal{S}} x^2 + y^2 + z^2 \, dS = \frac{R}{3} \int_{\mathcal{S}} dS = \frac{4\pi R^3}{3}$. Note that in each case the flux was calculated without ever having to start in on a parameterisation. Moral — look for symmetry tricks. The divergences in the three cases are 3, 0, 1 and it is easy to confirm that each flux integral agrees with the corresponding volume integral given by the divergence theorem.

A3 The divergence of ∇f is $12(x^2 + y^2 + z^2)$. (a) 12 (b) $48\pi R^5/5$ (use polar coordinates).

A4 (a) $(0, 0, 2)$ (b) $(0, 0, 0)$ (c) $(2x, -2y, 1)$.

A6 $\text{curl}(v) = 1$ so that

$$\text{Area}(\Omega) = \int_{\Omega} \text{Curl}(v) \, dA = \int_{\partial\Omega} v \cdot \hat{T} \, ds = \int_0^{2\pi} (0, \alpha \cos t) \cdot (-\alpha \sin t, \beta \cos t) \, dt = \pi\alpha\beta.$$

A7 One possibility is to parametrise \mathcal{S} by $x(u, v) = (u, v, 2 - u^2 - v^2)$ for $(u, v) \in \Omega = \{u^2 + v^2 \leq 2\}$. Then inward (non-unit) normal vector $\hat{N}(u, v) = (-2u, -2v, -1)$. By parameterising $\partial\mathcal{S}$ by $\alpha(t) = (\sqrt{2} \cos t, \sqrt{2} \sin t, 0)$ for $t \in [0, 2\pi]$ the tangent vector and normal vector are then suitably oriented for Stokes' theorem. Both sides of Stokes' identity give the value 2π , for example $\int_{\mathcal{S}} \nabla \times v \cdot \hat{N} \, dS = \int_{\Omega} (2u + 2v + 1) \, dudv = \int_{\Omega} dudv = \text{Area of } \Omega$, (using symmetry to reduce the integrand from $2u + 2v + 1$ to 1).

B1 (a) Summing the single parts (as in the lecture), result is $\frac{3}{2}$. (b) A possible parametrisation for \mathcal{S} is $\alpha(s, t) = (s \cos t, s \sin t, s^2)$ for $s \in [0, 1]$, $t \in [0, 2\pi]$. The unit normal in positive z -direction is then $\hat{N} = \frac{1}{\sqrt{4s^2+1}}(-2s \cos t, -2s \sin t, 1)$. Result: $-\pi/2$.

B2 (a) A simple calculation gives $\text{div } f = 1$, integrating this over the disc with radius R gives the result πR^2 . (b) Divergence = 4. Flux = $4 \times$ Volume of the pyramid with base area 9 and height 2 = 24.

B3 (a) Use the definitions of the cross product, of the curl and diligence. (b) Use the product rule (c) Chain rule.

B4 The expression for $\text{div}(fv)$ follows from the product rule, the integration by parts formula can be derived by applying Gauss's theorem. Adding IBP for f, g and g, f gives Green's identity.

C3 (a) Stokes theorem for the vector field $\underline{v} = (f, 0, 0)$ becomes

$$\int_C f \hat{T}_1 \, ds = \int_S \left(\frac{\partial f}{\partial z} N_2 - \frac{\partial f}{\partial y} N_3 \right) \, dS.$$

Now repeat with $\underline{v} = (0, f, 0)$ and $\underline{v} = (0, 0, f)$. (b) Prove each co-ordinate of the identity separately, each is a case of the divergence theorem.

C4 Since f_1 and f_2 vanish on $\partial\Omega$, the boundary terms in Green's identity vanish and we get $\lambda_2 \int_{\Omega} f_1 f_2 = - \int_{\Omega} f_1 \Delta f_2 = - \int_{\Omega} f_2 \Delta f_1 = \lambda_1 \int_{\Omega} f_1 f_2$. Since $\lambda_1 \neq \lambda_2$ this implies the result.