

# Computing Kleinian modular forms

Aurel Page

June 4, 2014

Warwick, LMFDB workshop

- 1 Motivation
- 2 Arithmetic Kleinian groups
- 3 Algorithms
- 4 Examples

# Elliptic curves and automorphic forms

Want a database of elliptic curves over number field  $F$ ?

# Elliptic curves and automorphic forms

Want a database of elliptic curves over number field  $F$ ?  
Start with  $GL_2$  automorphic forms!

Where do you find automorphic forms?

# Elliptic curves and automorphic forms

Want a database of elliptic curves over number field  $F$ ?  
Start with  $GL_2$  automorphic forms!

Where do you find automorphic forms?  
In the cohomology of arithmetic groups!

Matsushima's formula:  $\Gamma$  discrete cocompact subgroup of  
connected Lie group  $G$ ,  $E$  representation of  $G$ .

$$H^i(\Gamma, E) \cong \bigoplus_{\pi \in \widehat{G}} \text{Hom}(\pi, L^2(\Gamma \backslash G)) \otimes H^i(\mathfrak{g}, \mathcal{K}; \pi \otimes E)$$

# Automorphic forms for $GL_2$

Where do you find  $GL_2$  automorphic forms?

# Automorphic forms for $GL_2$

Where do you find  $GL_2$  automorphic forms?

- $H^*(GL_2(\mathbb{Z}_F), E)$

# Automorphic forms for $GL_2$

Where do you find  $GL_2$  automorphic forms?

- $H^*(GL_2(\mathbb{Z}_F), E)$
- Jacquet–Langlands:  $H^*(\mathcal{O}^\times, E)$ ,  $\mathcal{O}$  order in a quaternion algebra



# Automorphic forms for $GL_2$

Where do you find  $GL_2$  automorphic forms?

- $H^*(GL_2(\mathbb{Z}_F), E)$
- Jacquet–Langlands:  $H^*(\mathcal{O}^\times, E)$ ,  $\mathcal{O}$  order in a quaternion algebra

**Kleinian** case:  $\mathcal{O}^\times \subset GL_2(\mathbb{C})$ ,  $H^1(\mathcal{O}^\times, E)$

# Automorphic forms for $GL_2$

Where do you find  $GL_2$  automorphic forms?

- $H^*(GL_2(\mathbb{Z}_F), E)$
- Jacquet–Langlands:  $H^*(\mathcal{O}^\times, E)$ ,  $\mathcal{O}$  order in a quaternion algebra

**Kleinian** case:  $\mathcal{O}^\times \subset GL_2(\mathbb{C})$ ,  $H^1(\mathcal{O}^\times, E)$

Call  $H^i(\mathcal{O}^\times, E)$  a space of **Kleinian modular forms**.

# Motivation for the Kleinian case

- Elliptic curves, Abelian varieties of  $GL_2$  type over number fields.

## Motivation for the Kleinian case

- Elliptic curves, Abelian varieties of  $GL_2$  type over number fields.
- Torsion phenomenon.

## Motivation for the Kleinian case

- Elliptic curves, Abelian varieties of  $GL_2$  type over number fields.
- Torsion phenomenon.
- Attached Galois representations: open case.

# Arithmetic Kleinian groups

# Kleinian groups

$\mathbb{H} := \mathbb{C} + \mathbb{C}j$  where  $j^2 = -1$  and  $jz = \bar{z}j$  for all  $z \in \mathbb{C}$ .

The upper half-space  $\mathcal{H}^3 := \mathbb{C} + \mathbb{R}_{>0}j$ .

Metric  $ds^2 = \frac{|dz|^2 + dt^2}{t^2}$ , volume  $dV = \frac{dx dy dt}{t^3}$ .

# Kleinian groups

$\mathbb{H} := \mathbb{C} + \mathbb{C}j$  where  $j^2 = -1$  and  $jz = \bar{z}j$  for all  $z \in \mathbb{C}$ .

The upper half-space  $\mathcal{H}^3 := \mathbb{C} + \mathbb{R}_{>0}j$ .

Metric  $ds^2 = \frac{|dz|^2 + dt^2}{t^2}$ , volume  $dV = \frac{dx dy dt}{t^3}$ .

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot w := (aw + b)(cw + d)^{-1} \text{ for } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C}).$$



# Kleinian groups

$\mathbb{H} := \mathbb{C} + \mathbb{C}j$  where  $j^2 = -1$  and  $jz = \bar{z}j$  for all  $z \in \mathbb{C}$ .

The upper half-space  $\mathcal{H}^3 := \mathbb{C} + \mathbb{R}_{>0}j$ .

Metric  $ds^2 = \frac{|dz|^2 + dt^2}{t^2}$ , volume  $dV = \frac{dx dy dt}{t^3}$ .

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot w := (aw + b)(cw + d)^{-1} \text{ for } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C}).$$

Kleinian group: discrete subgroup of  $\mathrm{SL}_2(\mathbb{C})$ .

Cofinite if it has finite covolume.

# Quaternion algebras

A quaternion algebra  $B$  over a field  $F$  is a central simple algebra of dimension 4 over  $F$ .

Explicitly,  $B = \left(\frac{a,b}{F}\right) = F + Fi + Fj + Fij$ ,  $i^2 = a$ ,  $j^2 = b$ ,  $ij = -ji$  (char  $F \neq 2$ ).

# Quaternion algebras

A quaternion algebra  $B$  over a field  $F$  is a central simple algebra of dimension 4 over  $F$ .

Explicitly,  $B = \left(\frac{a,b}{F}\right) = F + Fi + Fj + Fij$ ,  $i^2 = a$ ,  $j^2 = b$ ,  $ij = -ji$  (char  $F \neq 2$ ).

Reduced norm:  $\text{nrd}(x + yi + zj + tij) = x^2 - ay^2 - bz^2 + abt^2$ .

# Quaternion algebras

A quaternion algebra  $B$  over a field  $F$  is a central simple algebra of dimension 4 over  $F$ .

Explicitly,  $B = \left(\frac{a,b}{F}\right) = F + Fi + Fj + Fij$ ,  $i^2 = a$ ,  $j^2 = b$ ,  $ij = -ji$  (char  $F \neq 2$ ).

Reduced norm:  $\text{nrd}(x + yi + zj + tij) = x^2 - ay^2 - bz^2 + abt^2$ .

$\mathbb{Z}_F$  integers of  $F$  number field.

Order  $\mathcal{O} \subset B$ : subring, finitely generated  $\mathbb{Z}_F$ -module,  $\mathcal{O}F = B$ .

# Quaternion algebras

A quaternion algebra  $B$  over a field  $F$  is a central simple algebra of dimension 4 over  $F$ .

Explicitly,  $B = \left(\frac{a,b}{F}\right) = F + Fi + Fj + Fij$ ,  $i^2 = a$ ,  $j^2 = b$ ,  $ij = -ji$  (char  $F \neq 2$ ).

Reduced norm:  $\text{nrd}(x + yi + zj + tij) = x^2 - ay^2 - bz^2 + abt^2$ .

$\mathbb{Z}_F$  integers of  $F$  number field.

Order  $\mathcal{O} \subset B$ : subring, finitely generated  $\mathbb{Z}_F$ -module,  $\mathcal{O}F = B$ .

A place  $v$  of  $F$  is split or ramified according as whether  $B \otimes_F F_v$  is  $\mathcal{M}_2(F_v)$  or a division algebra.

## Covolume formula

$F$  almost totally real number field: exactly one complex place.

$B/F$  **Kleinian** quaternion algebra: ramified at every real place.

## Covolume formula

$F$  almost totally real number field: exactly one complex place.

$B/F$  Kleinian quaternion algebra: ramified at every real place.

$\mathcal{O}$  order in  $B$ ,  $\Gamma$  image of reduced norm one group  $\mathcal{O}^1$  in  $\mathrm{SL}_2(\mathbb{C})$ .

## Covolume formula

$F$  almost totally real number field: exactly one complex place.

$B/F$  **Kleinian** quaternion algebra: ramified at every real place.

$\mathcal{O}$  order in  $B$ ,  $\Gamma$  image of reduced norm one group  $\mathcal{O}^1$  in  $SL_2(\mathbb{C})$ .

### Theorem

$\Gamma$  is a cofinite Kleinian group.

It is cocompact iff  $B$  is a division algebra.

If  $\mathcal{O}$  is maximal then

$$\text{Covol}(\Gamma) = \frac{|\Delta_F|^{3/2} \zeta_F(2) \prod_{\mathfrak{p} \text{ ram.}} (N(\mathfrak{p}) - 1)}{(4\pi^2)^{[F:\mathbb{Q}] - 1}}.$$



# Algorithms

# Sketch of algorithm

- 1 Compute  $B$  and  $\mathcal{O}$

## Sketch of algorithm

- 1 Compute  $B$  and  $\mathcal{O}$
- 2 Compute  $\mathcal{O}^1$

## Sketch of algorithm

- 1 Compute  $B$  and  $\mathcal{O}$
- 2 Compute  $\mathcal{O}^1$
- 3 Compute  $H^1(\mathcal{O}^1)$

## Sketch of algorithm

- 1 Compute  $B$  and  $\mathcal{O}$
- 2 Compute  $\mathcal{O}^1$
- 3 Compute  $H^1(\mathcal{O}^1)$
- 4 Compute generator  $\delta$  of ideal of norm  $\mathfrak{p}$

## Sketch of algorithm

- 1 Compute  $B$  and  $\mathcal{O}$
- 2 Compute  $\mathcal{O}^1$
- 3 Compute  $H^1(\mathcal{O}^1)$
- 4 Compute generator  $\delta$  of ideal of norm  $\mathfrak{p}$
- 5 Compute Hecke operator  $T_\delta$  on  $H^1(\mathcal{O}^1)$

# Fundamental domains

$\Gamma$  a Kleinian group. An open subset  $\mathcal{F} \subset \mathcal{H}^3$  is a fundamental domain if

- $\Gamma \cdot \overline{\mathcal{F}} = \mathcal{H}^3$
- $\mathcal{F} \cap \gamma\mathcal{F} = \emptyset$  for all  $1 \neq \gamma \in \Gamma$ .

# Dirichlet domains



# Dirichlet domains

Let  $p \in \mathcal{H}^3$  with trivial stabilizer in  $\Gamma$ . The Dirichlet domain

$$\begin{aligned} D_p(\Gamma) &:= \{x \in X \mid d(x, p) < d(\gamma \cdot x, p) \forall \gamma \in \Gamma \setminus \{1\}\} \\ &= \{x \in X \mid d(x, p) < d(x, \gamma^{-1} \cdot p) \forall \gamma \in \Gamma \setminus \{1\}\}. \end{aligned}$$

# Dirichlet domains

Let  $p \in \mathcal{H}^3$  with trivial stabilizer in  $\Gamma$ . The Dirichlet domain

$$\begin{aligned} D_p(\Gamma) &:= \{x \in X \mid d(x, p) < d(\gamma \cdot x, p) \forall \gamma \in \Gamma \setminus \{1\}\} \\ &= \{x \in X \mid d(x, p) < d(x, \gamma^{-1} \cdot p) \forall \gamma \in \Gamma \setminus \{1\}\}. \end{aligned}$$

is a fundamental domain for  $\Gamma$  that is a hyperbolic polyhedron. If  $\Gamma$  is cofinite,  $D_p(\Gamma)$  has finitely many faces.

# Structure of the Dirichlet domain

## Structure of the Dirichlet domain

- Faces of the Dirichlet domain are grouped into **pairs**, corresponding to elements  $g, g^{-1} \in \Gamma$ ;

## Structure of the Dirichlet domain

- Faces of the Dirichlet domain are grouped into **pairs**, corresponding to elements  $g, g^{-1} \in \Gamma$ ;
- Edges of the domain are grouped into **cycles**, product of corresponding elements in  $\Gamma$  has **finite order**.

# Poincaré's theorem

## Theorem (Poincaré)

- *The elements corresponding to the faces are generators of  $\Gamma$ . The relations corresponding to the edge cycles generate all the relations among the generators.*

# Poincaré's theorem

## Theorem (Poincaré)

- *The elements corresponding to the faces are generators of  $\Gamma$ . The relations corresponding to the edge cycles generate all the relations among the generators.*
- *If a partial Dirichlet domain  $D_p(S)$  has a face-pairing and cycles of edges, then it is a fundamental domain for the group generated by  $S$ .*

## Reduction algorithm

$x \in \mathcal{H}^3$ ,  $S \subset \Gamma$  corresponding to the faces of  $D_p(\Gamma)$ .



## Reduction algorithm

$x \in \mathcal{H}^3$ ,  $S \subset \Gamma$  corresponding to the faces of  $D_p(\Gamma)$ .

- if possible  $x \leftarrow gx$  for some  $g \in S$  s.t.  $d(x, p)$  decreases

## Reduction algorithm

$x \in \mathcal{H}^3$ ,  $S \subset \Gamma$  corresponding to the faces of  $D_p(\Gamma)$ .

- if possible  $x \leftarrow gx$  for some  $g \in S$  s.t.  $d(x, p)$  decreases
- repeat

## Reduction algorithm

$x \in \mathcal{H}^3$ ,  $S \subset \Gamma$  corresponding to the faces of  $D_p(\Gamma)$ .

- if possible  $x \leftarrow gx$  for some  $g \in S$  s.t.  $d(x, p)$  decreases
- repeat

$\rightarrow$  point  $x' = g_k \cdots g_1 x$  s.t.  $x' \in D_p(\Gamma)$ .

## Reduction algorithm

$x \in \mathcal{H}^3$ ,  $S \subset \Gamma$  corresponding to the faces of  $D_p(\Gamma)$ .

- if possible  $x \leftarrow gx$  for some  $g \in S$  s.t.  $d(x, p)$  decreases
- repeat

$\rightarrow$  point  $x' = g_k \cdots g_1 x$  s.t.  $x' \in D_p(\Gamma)$ .

In particular for  $\gamma \in \Gamma$ , take  $x = \gamma^{-1}p$ , which will reduce to  $x' = p$ , to write  $\gamma$  as a product of the generators.

# Computation in hyperbolic space

# Computation in hyperbolic space

We have :

- an explicit formula for the hyperbolic distance;

# Computation in hyperbolic space

We have :

- an explicit formula for the hyperbolic distance;
- an explicit formula for bisectors;

# Computation in hyperbolic space

We have :

- an explicit formula for the hyperbolic distance;
- an explicit formula for bisectors;
- an algorithm for computing intersections of half-spaces;



# Computation in hyperbolic space

We have :

- an explicit formula for the hyperbolic distance;
- an explicit formula for bisectors;
- an algorithm for computing intersections of half-spaces;
- an algorithm for computing the volume of a polyhedron.

# Algorithm

Suppose  $\mathcal{O}$  maximal.

# Algorithm

Suppose  $\mathcal{O}$  maximal.

- Enumerate elements of  $\mathcal{O}^1$  in a finite set  $S$

# Algorithm

Suppose  $\mathcal{O}$  maximal.

- Enumerate elements of  $\mathcal{O}^1$  in a finite set  $S$
- Compute  $D = D_p(S)$

# Algorithm

Suppose  $\mathcal{O}$  maximal.

- Enumerate elements of  $\mathcal{O}^1$  in a finite set  $S$
- Compute  $D = D_p(S)$
- Repeat until  $D$  has a face-pairing and  $\text{Vol}(D) < 2 \text{Covol}(\Gamma)$

# Complexity

Why would you care about the complexity?

# Complexity

Why would you care about the complexity?

Bound on the size of the Dirichlet domain:

# Complexity

Why would you care about the complexity?

Bound on the size of the Dirichlet domain:

$\mathrm{SL}_2(\mathbb{C})$  action on  $L_0^2(\Gamma \backslash \mathrm{SL}_2(\mathbb{C}))$  mixes things fast.



# Complexity

Why would you care about the complexity?

Bound on the size of the Dirichlet domain:

$\mathrm{SL}_2(\mathbb{C})$  action on  $L_0^2(\Gamma \backslash \mathrm{SL}_2(\mathbb{C}))$  mixes things fast.  
 $\Rightarrow \mathrm{diam}(D_\rho(\Gamma)) \ll \log \mathrm{Covol}(\Gamma)$

# Complexity

Why would you care about the complexity?

Bound on the size of the Dirichlet domain:

$\mathrm{SL}_2(\mathbb{C})$  action on  $L_0^2(\Gamma \backslash \mathrm{SL}_2(\mathbb{C}))$  mixes things fast.

$\Rightarrow \mathrm{diam}(D_\rho(\Gamma)) \ll \log \mathrm{Covol}(\Gamma)$

- Proved complexity:  $\mathrm{Covol}(\Gamma)^{O(1)}$
- Observed complexity:  $\mathrm{Covol}(\Gamma)^2$
- Lower bound:  $\mathrm{Covol}(\Gamma)$

# Group cohomology

# Group cohomology

Cocycles :

$$Z^1(\Gamma, E) := \{\phi : \Gamma \rightarrow E \mid \phi(gh) = \phi(g) + g \cdot \phi(h) \forall g, h \in \Gamma\}$$

# Group cohomology

Cocycles :

$$Z^1(\Gamma, E) := \{\phi : \Gamma \rightarrow E \mid \phi(gh) = \phi(g) + g \cdot \phi(h) \forall g, h \in \Gamma\}$$

Coboundaries :

$$B^1(\Gamma, E) := \{\phi_x : g \mapsto x - g \cdot x : x \in E\} \subset Z^1(\Gamma, E)$$

# Group cohomology

Cocycles :

$$Z^1(\Gamma, E) := \{\phi : \Gamma \rightarrow E \mid \phi(gh) = \phi(g) + g \cdot \phi(h) \forall g, h \in \Gamma\}$$

Coboundaries :

$$B^1(\Gamma, E) := \{\phi_x : g \mapsto x - g \cdot x : x \in E\} \subset Z^1(\Gamma, E)$$

Cohomology :

$$H^1(\Gamma, E) := Z^1(\Gamma, E)/B^1(\Gamma, E)$$

# Hecke operators

Let  $\delta \in \text{Comm}(\Gamma)$ .

$$H^i(\Gamma, M)$$

# Hecke operators

Let  $\delta \in \text{Comm}(\Gamma)$ .

$$\begin{array}{c}
 H^i(\Gamma, M) \\
 \text{Res} \downarrow \\
 H^i(\Gamma \cap \delta\Gamma\delta^{-1}, M)
 \end{array}$$



# Hecke operators

Let  $\delta \in \text{Comm}(\Gamma)$ .

$$\begin{array}{ccc}
 H^i(\Gamma, M) & & \\
 \text{Res} \downarrow & & \\
 H^i(\Gamma \cap \delta\Gamma\delta^{-1}, M) & \xrightarrow{\tilde{\delta}} & H^i(\delta^{-1}\Gamma\delta \cap \Gamma, M)
 \end{array}$$

# Hecke operators

Let  $\delta \in \text{Comm}(\Gamma)$ .

$$\begin{array}{ccc}
 H^i(\Gamma, M) & & H^i(\Gamma, M) \\
 \text{Res} \downarrow & & \uparrow \text{Cores} \\
 H^i(\Gamma \cap \delta\Gamma\delta^{-1}, M) & \xrightarrow{\tilde{\delta}} & H^i(\delta^{-1}\Gamma\delta \cap \Gamma, M)
 \end{array}$$

# Hecke operators

Let  $\delta \in \text{Comm}(\Gamma)$ .

$$\begin{array}{ccc}
 H^i(\Gamma, M) & \xrightarrow{T_\delta} & H^i(\Gamma, M) \\
 \text{Res} \downarrow & & \uparrow \text{Cores} \\
 H^i(\Gamma \cap \delta\Gamma\delta^{-1}, M) & \xrightarrow{\tilde{\delta}} & H^i(\delta^{-1}\Gamma\delta \cap \Gamma, M)
 \end{array}$$

# Principal ideal problem

Recall: have to find a generator  $\delta$  of ideal of norm  $\mathfrak{p} = (\pi)$ .

# Principal ideal problem

Recall: have to find a generator  $\delta$  of ideal of norm  $\mathfrak{p} = (\pi)$ .

Search in  $\mathcal{O}^1 \setminus \{\text{nrd} = \pi\}$ : complexity  $\Delta_B^{O(1)}$ .

# Principal ideal problem

Recall: have to find a generator  $\delta$  of ideal of norm  $\mathfrak{p} = (\pi)$ .

Search in  $\mathcal{O}^1 \setminus \{\text{nrd} = \pi\}$ : complexity  $\Delta_B^{O(1)}$ .

Buchmann's algorithm over number fields: subexponential under GRH.

# Principal ideal problem

Recall: have to find a generator  $\delta$  of ideal of norm  $\mathfrak{p} = (\pi)$ .

Search in  $\mathcal{O}^1 \setminus \{\text{nrd} = \pi\}$ : complexity  $\Delta_B^{O(1)}$ .

Buchmann's algorithm over number fields: subexponential under GRH.

Adapt Buchmann's algorithm over a quaternion algebra: heuristically subexponential.

# Examples



## A quartic example

Let  $F$  the unique quartic field of signature  $(2, 1)$  and discriminant  $-275$ . Let  $B$  be the unique quaternion algebra with discriminant  $11\mathbb{Z}_F$ , ramified at every real place of  $F$ . Let  $\mathcal{O}$  be a maximal order in  $B$  (it is unique up to conjugation). Then  $\mathcal{O}^1$  is a Kleinian group with covolume  $93.72\dots$ . The fundamental domain I have computed has 310 faces and 924 edges.

## A quartic example

Let  $F$  the unique quartic field of signature  $(2, 1)$  and discriminant  $-275$ . Let  $B$  be the unique quaternion algebra with discriminant  $11\mathbb{Z}_F$ , ramified at every real place of  $F$ . Let  $\mathcal{O}$  be a maximal order in  $B$  (it is unique up to conjugation). Then  $\mathcal{O}^1$  is a Kleinian group with covolume  $93.72\dots$ . The fundamental domain I have computed has 310 faces and 924 edges.

Have a look at the fundamental domain!

# A quartic example

$N(p)$	$T_p$	characteristic polynomial
9	$\begin{pmatrix} 2 & -4 & -5 \\ -2 & 3 & -2 \\ 0 & 0 & -5 \end{pmatrix}$	$(x + 5)(x^2 - 5x - 2)$
9	same	$(x + 5)(x^2 - 5x - 2)$
16	$\begin{pmatrix} 1 & 4 & -11 \\ 2 & 0 & -6 \\ 0 & 0 & -8 \end{pmatrix}$	$(x + 8)(x^2 - x - 8)$
19	$\begin{pmatrix} -4 & 0 & 4 \\ 0 & -4 & 2 \\ 0 & 0 & 0 \end{pmatrix}$	$x(x + 4)^2$
19	same	$x(x + 4)^2$
25	—	$(x + 9)(x^2 - x - 74)$
29	—	$x(x^2 + 6x - 24)$
29	—	$x(x^2 + 6x - 24)$

## Non base-change classes

Bianchi groups:  $\Gamma = \mathrm{SL}_2(\mathbb{Z}_F)$  with  $F$  quadratic imaginary.

## Non base-change classes

Bianchi groups:  $\Gamma = \mathrm{SL}_2(\mathbb{Z}_F)$  with  $F$  quadratic imaginary.  
Finis, Grunewald, Tiraó: formula for the base-change subspace  
in  $H^i(\Gamma, E_k)$ . Remark: very few non base-change classes.

## Non base-change classes

Bianchi groups:  $\Gamma = \mathrm{SL}_2(\mathbb{Z}_F)$  with  $F$  quadratic imaginary.  
Finis, Grunewald, Tiraó: formula for the base-change subspace  
in  $H^i(\Gamma, E_k)$ . Remark: very few non base-change classes.  
Şengün, Rahm: extended their computations.

## Non base-change classes

Bianchi groups:  $\Gamma = \mathrm{SL}_2(\mathbb{Z}_F)$  with  $F$  quadratic imaginary.  
Finis, Grunewald, Tirao: formula for the base-change subspace in  $H^i(\Gamma, E_k)$ . Remark: very few non base-change classes.  
Şengün, Rahm: extended their computations.

### Conjecture (Modified Maeda's conjecture)

*The set of non-lifted cuspidal newforms in  $H^i(\Gamma, E_k)$ , modulo twins, forms one Galois orbit.*

## Non base-change classes

Bianchi groups:  $\Gamma = \mathrm{SL}_2(\mathbb{Z}_F)$  with  $F$  quadratic imaginary.  
 Finis, Grunewald, Tiraó: formula for the base-change subspace  
 in  $H^i(\Gamma, E_k)$ . Remark: very few non base-change classes.  
 Şengün, Rahm: extended their computations.

### Conjecture (Modified Maeda's conjecture)

*The set of non-lifted cuspidal newforms in  $H^i(\Gamma, E_k)$ , modulo  
 twins, forms one Galois orbit.*

Experiment : always dimension 2 space,  
 except  $(d, k) = (-199, 2)$ : dimension 4.



## Non base-change classes

Bianchi groups:  $\Gamma = \mathrm{SL}_2(\mathbb{Z}_F)$  with  $F$  quadratic imaginary.  
 Finis, Grunewald, Tirao: formula for the base-change subspace  
 in  $H^i(\Gamma, E_k)$ . Remark: very few non base-change classes.  
 Şengün, Rahm: extended their computations.

### Conjecture (Modified Maeda's conjecture)

*The set of non-lifted cuspidal newforms in  $H^i(\Gamma, E_k)$ , modulo  
 twins, forms one Galois orbit.*

Experiment : always dimension 2 space,  
 except  $(d, k) = (-199, 2)$ : dimension 4. Two twin Galois orbits  
 with coefficients in  $\mathbb{Q}(\sqrt{13})$ , swapped by  $\mathrm{Gal}(F/\mathbb{Q})$ .

# Everywhere good reduction abelian surfaces

Bianchi group for  $F = \mathbb{Q}(\sqrt{-223})$ .

## Everywhere good reduction abelian surfaces

Bianchi group for  $F = \mathbb{Q}(\sqrt{-223})$ .

Have a look at the fundamental domain!

## Everywhere good reduction abelian surfaces

Bianchi group for  $F = \mathbb{Q}(\sqrt{-223})$ .

Have a look at the fundamental domain!

Cuspidal subspace in  $H^1(\Gamma, \mathbb{C})$ : dimension 2, one Galois orbit in  $\mathbb{Q}(\sqrt{5})$ .

## Everywhere good reduction abelian surfaces

Bianchi group for  $F = \mathbb{Q}(\sqrt{-223})$ .

Have a look at the fundamental domain!

Cuspidal subspace in  $H^1(\Gamma, \mathbb{C})$ : dimension 2, one Galois orbit in  $\mathbb{Q}(\sqrt{5})$ .

Şengün, Dembélé : the Jacobian  $J$  of the hyperelliptic curve

$$y^2 = 33x^6 + 110\sqrt{-223}x^5 + 36187x^4 - 28402\sqrt{-223}x^3 - 2788739x^2 + 652936\sqrt{-223}x + 14157596$$

has good reduction everywhere,  $\text{End}(J) \otimes \mathbb{Q} \cong \mathbb{Q}(\sqrt{5})$  and matches the Hecke eigenvalues.

## Everywhere good reduction abelian surfaces

Bianchi group for  $F = \mathbb{Q}(\sqrt{-223})$ .

Have a look at the fundamental domain!

Cuspidal subspace in  $H^1(\Gamma, \mathbb{C})$ : dimension 2, one Galois orbit in  $\mathbb{Q}(\sqrt{5})$ .

Şengün, Dembélé : the Jacobian  $J$  of the hyperelliptic curve

$$y^2 = 33x^6 + 110\sqrt{-223}x^5 + 36187x^4 - 28402\sqrt{-223}x^3 - 2788739x^2 + 652936\sqrt{-223}x + 14157596$$

has good reduction everywhere,  $\text{End}(J) \otimes \mathbb{Q} \cong \mathbb{Q}(\sqrt{5})$  and matches the Hecke eigenvalues.

Looking for more examples in  $\mathbb{Q}(\sqrt{-455})$  and  $\mathbb{Q}(\sqrt{-571})$ .

# Torsion Jacquet-Langlands

Can also look at  $H_1(\Gamma, \mathbb{Z})_{\text{tor}}$  and  $H_1(\Gamma, \mathbb{F}_p)$ .

## Torsion Jacquet-Langlands

Can also look at  $H_1(\Gamma, \mathbb{Z})_{\text{tor}}$  and  $H_1(\Gamma, \mathbb{F}_p)$ .

Jacquet–Langlands: relation between (co)homology of  $\Gamma_0(\mathfrak{N})$  in a Bianchi group and that of  $\Gamma$  coming from a division algebra of discriminant  $\mathfrak{N}$ .



# Torsion Jacquet-Langlands

Can also look at  $H_1(\Gamma, \mathbb{Z})_{\text{tor}}$  and  $H_1(\Gamma, \mathbb{F}_p)$ .

Jacquet–Langlands: relation between (co)homology of  $\Gamma_0(\mathfrak{N})$  in a Bianchi group and that of  $\Gamma$  coming from a division algebra of discriminant  $\mathfrak{N}$ .

Calegari, Venkatesh: numerical torsion Jacquet-Langlands correspondence.

## Torsion Jacquet-Langlands

Can also look at  $H_1(\Gamma, \mathbb{Z})_{\text{tor}}$  and  $H_1(\Gamma, \mathbb{F}_p)$ .

Jacquet–Langlands: relation between (co)homology of  $\Gamma_0(\mathfrak{N})$  in a Bianchi group and that of  $\Gamma$  coming from a division algebra of discriminant  $\mathfrak{N}$ .

Calegari, Venkatesh: numerical torsion Jacquet-Langlands correspondence.

Joint with H. Şengün: experimental verification with Hecke operators (still in progress).

Thank you for your attention !