



University of
Zurich ^{UZH}



Critical phases for non-relativistic 2d interacting bosons: Renormalization Group results


Serena Cenatiempo

joint work with A. Giuliani, Università di Roma Tre


Many-Body Quantum Systems
University of Warwick

March 20, 2014


Ultracolds atoms: an ultralow temperature laboratory for many-body physics




Hänsel et al., "Atom chip" (2001)





Krüger et al. (2007)



Billy et al., Guided atom laser (2008)



Nature Physics (2005)



Greiner et al., Quantum Phase Transition (2002)

The model

- ▶ Gas of non-relativistic bosons in a periodic box $\Omega_L \in \mathbb{R}^d$, $|\Omega_L| = L^d$
- ▶ weak repulsive short range two-body potential $\lambda v(\vec{x})$, $0 < \lambda \ll 1$

$$H_{N,L} = - \sum_{i=1}^N \Delta_{\vec{x}_i} + \lambda \sum_{1 \leq i < j \leq N} v(\vec{x}_i - \vec{x}_j) \quad \text{on } \mathcal{H}_N^{\text{sym}}$$

The model

- ▶ Gas of non-relativistic bosons in a periodic box $\Omega_L \in \mathbb{R}^d$, $|\Omega_L| = L^d$
- ▶ weak repulsive short range two-body potential $\lambda v(\vec{x})$, $0 < \lambda \ll 1$

$$H_{N,L} = - \sum_{i=1}^N \Delta_{\vec{x}_i} + \lambda \sum_{1 \leq i < j \leq N} v(\vec{x}_i - \vec{x}_j) \quad \text{on } \mathcal{H}_N^{\text{sym}}$$

Goal: to construct the thermal ground state in infinite volume

$$e_0(\rho) = - \lim_{\beta \rightarrow \infty} \lim_{L \rightarrow \infty} \frac{1}{\beta L^d} \log \text{Tr}_{\mathcal{H}_N^{\text{sym}}} e^{-\beta(H_{N,L} - \mu_{\beta,L} N)}$$

$$S(\vec{x}, \vec{y}) = \lim_{\beta \rightarrow \infty} \lim_{L \rightarrow \infty} \frac{\text{Tr}_{\mathcal{H}_N^{\text{sym}}} [e^{-\beta(H_{N,L} - \mu_{\beta,L} N)} a_{\vec{x}}^+ a_{\vec{y}}]}{\text{Tr}_{\mathcal{H}_N^{\text{sym}}} e^{-\beta(H_{N,L} - \mu_{\beta,L} N)}}$$

with $\mu_{\beta,L} \leq 0$ the chemical potential.

Existence of condensation

The non-interacting case (Einstein, 1925)

	$\lambda = 0$		
$d = 3$	$0 \leq T \leq T_c^0$		
$d = 2$	$T=0$		
$d = 1$	$T=0$		

The density of the states with $\vec{k} \neq 0$ is bounded in each dimension at $T = 0$ and at finite T in $3d$ as $\mu_{L,\beta} \rightarrow 0$:

$$\lim_{|\Omega| \rightarrow +\infty} \rho_{\Omega,\beta}^{(\vec{k} \neq 0)} = \int \frac{d^d \vec{k}}{(2\pi)^d} \frac{1}{e^{\beta(\vec{k}^2 - \mu_{L,\beta})} - 1} \leq \rho_{\beta}^{\text{critical}}$$

To fix the system at $\rho = \rho_0 + \rho_{\beta}^{\text{critical}}$ the chemical potential has to be chosen such that $\lim_{\beta \rightarrow \infty} \lim_{L \rightarrow \infty} (-\mu_{\beta,L}) = 1/\rho_0$.

Existence of condensation

The interacting case within Bogoliubov approximation

	$\lambda = 0$	$\lambda > 0$ & Bogoliubov	
$d = 3$	$0 \leq T \leq T_c^0$	$0 \leq T \leq T_c^B$	
$d = 2$	$T=0$	$T=0$	
$d = 1$	$T=0$	No cond.	

Total density according to Bogoliubov approximation

$$\rho = \rho_0 + \underbrace{\int \frac{d^d \vec{k}}{(2\pi)^d} \frac{F(\vec{k}) - \varepsilon(\vec{k})}{\varepsilon(\vec{k})}}_{\substack{\approx \\ \vec{k} \approx 0}} + \underbrace{\int \frac{d^d \vec{k}}{(2\pi)^d} \frac{F(\vec{k})}{\varepsilon(\vec{k})} \frac{1}{e^{\beta \varepsilon(\vec{k})} - 1}}_{\substack{\approx \\ \vec{k} \approx 0, \beta \text{ finite}}} \frac{1}{\beta |\vec{k}|^2}$$

with $F(\vec{k}) = |\vec{k}|^2 + \lambda \hat{v}(\vec{k}) \rho_0$ and $\varepsilon^2(\vec{k}) = |\vec{k}|^4 + 2 |\vec{k}|^2 \lambda \hat{v}(\vec{k}) \rho_0$.

Existence of condensation

The interacting case: known results on condensation

	$\lambda = 0$	$\lambda > 0$ & Bogoliubov	$\lambda > 0$
$d = 3$	$0 \leq T \leq T_c^0$	$0 \leq T \leq T_c^B$	$0 \leq T \leq T_c ?$
$d = 2$	$T=0$	$T=0$	$T=0 ?$
$d = 1$	$T=0$	No cond.	No cond.

- ▶ Hard-core 3d bosons on a lattice at half filling
(Dyson, Lieb and Simon, 1978)

Existence of condensation

The interacting case: known results on condensation

	$\lambda = 0$	$\lambda > 0$ & Bogoliubov	$\lambda > 0$
$d = 3$	$0 \leq T \leq T_c^0$	$0 \leq T \leq T_c^B$	$0 \leq T \leq T_c ?$
$d = 2$	$T=0$	$T=0$	$T=0 ?$
$d = 1$	$T=0$	No cond.	No cond. ⁽²⁾

} MWH
thm⁽¹⁾

- Hard-core 3d bosons on a lattice at half filling
(Dyson, Lieb and Simon, 1978)

⁽¹⁾Hohenberg (1967) ⁽²⁾Lieb and Liniger (1963)

Existence of condensation

The interacting case: known results on condensation

	$\lambda = 0$	$\lambda > 0$ & Bogoliubov	$\lambda > 0$
$d = 3$	$0 \leq T \leq T_c^0$	$0 \leq T \leq T_c^B$	$0 \leq T \leq T_c ?$
$d = 2$	$T=0$	$T=0$	$T=0 ?$
$d = 1$	$T=0$	No cond.	No cond. ⁽²⁾

} MWH thm⁽¹⁾

- ▶ Hard-core 3d bosons on a lattice at half filling (Dyson, Lieb and Simon, 1978)
- ▶ 3d and 2d bosons in the Gross–Pitaevskii limit: $N/L = (\text{const.})$ (Lieb, Seiringer, Yngvason 2002)
- ▶ Bogoliubov's scheme has been proved to be valid in the mean field regime (Seiringer 2010, Dereziński and Napiórkowski 2013)

⁽¹⁾Hohenberg (1967) ⁽²⁾Lieb and Liniger (1963)

Existence of condensation

The interacting case: which general results for homogeneous bosons?

	$\lambda = 0$	$\lambda > 0$ & Bogoliubov	$\lambda > 0$
$d = 3$	$0 \leq T \leq T_c^0$	$0 \leq T \leq T_c^B$	$0 \leq T \leq T_c ?$
$d = 2$	$T=0$	$T=0$	$T=0 ?$
$d = 1$	$T=0$	No cond.	No cond.

Existence of condensation

The interacting case: which general results for homogeneous bosons?

	$\lambda = 0$	$\lambda > 0$ & Bogoliubov	$\lambda > 0$
$d = 3$	$0 \leq T \leq T_c^0$	$0 \leq T \leq T_c^B$	$0 \leq T \leq T_c ?$
$d = 2$	$T=0$	$T=0$	$T=0 ?$
$d = 1$	$T=0$	No cond.	No cond.

Corrections to Bogoliubov's via perturbation theory: Beliaev (1958), Hugenholtz & Pines (1959), Lee & Yang (1960), Gavoret & Nozières (1964), Nepomnyashchy & Nepomnyashchy (1978), Popov (1987).

Existence of condensation

The interacting case: which general results for homogeneous bosons?

	$\lambda = 0$	$\lambda > 0$ & Bogoliubov	$\lambda > 0$
$d = 3$	$0 \leq T \leq T_c^0$	$0 \leq T \leq T_c^B$	$0 \leq T \leq T_c ?$
$d = 2$	$T=0$	$T=0$	$T=0 ?$
$d = 1$	$T=0$	No cond.	No cond.

Corrections to Bogoliubov's via perturbation theory: Beliaev (1958), Hugenholtz & Pines (1959), Lee & Yang (1960), Gavoret & Nozières (1964), Nepomnyashchy & Nepomnyashchy (1978), Popov (1987).

Benfatto (1994 & 1997): proof of the order by order convergence in renormalized theory for the 3d case at $T = 0$ & strong BEC justification.

Existence of condensation

The interacting case: which general results for homogeneous bosons?

	$\lambda = 0$	$\lambda > 0$ & Bogoliubov	$\lambda > 0$
$d = 3$	$0 \leq T \leq T_c^0$	$0 \leq T \leq T_c^B$	$0 \leq T \leq T_c ?$
$d = 2$	$T=0$	$T=0$	$T=0 ?$
$d = 1$	$T=0$	No cond.	No cond.

Corrections to Bogoliubov's via perturbation theory: Beliaev (1958), Hugenholtz & Pines (1959), Lee & Yang (1960), Gavoret & Nozières (1964), Nepomnyashchy & Nepomnyashchy (1978), Popov (1987).

Benfatto (1994 & 1997): proof of the order by order convergence in renormalized theory for the 3d case at $T = 0$ & strong BEC justification.

Constructing the theory: a great challenge. A program addressing this issue started by Balaban, Feldman, Knörrer, Trubowitz (2008–2010)

Two dimensions

- Pistoiesi, Castellani, Di Castro, Strinati (1997 & 2004):
RG analysis in $3d$ and $2d$ by using **local Ward identities**
in a **dimensional regularization scheme** with $d = 3 - \varepsilon$.

After assuming the existence of a $O(1)$ fixed point
(suggested by a one loop calculation) they find that:

- * the condensate is stable in $2d$ and $T = 0$;
- * the theory displays a linear excitation spectrum a'la Bogoliubov.

Two dimensions

- ▶ Pistoiesi, Castellani, Di Castro, Strinati (1997 & 2004):
RG analysis in $3d$ and $2d$ by using **local Ward identities**
in a **dimensional regularization scheme** with $d = 3 - \varepsilon$.

After assuming the existence of a $O(1)$ fixed point
(suggested by a one loop calculation) they find that:

- * the condensate is stable in $2d$ and $T = 0$;
- * the theory displays a linear excitation spectrum a'la Bogoliubov.

- ▶ **Problems:**

- (1) Not even order by order results in $2d$ were available

Two dimensions

- Pistoiesi, Castellani, Di Castro, Strinati (1997 & 2004):
RG analysis in $3d$ and $2d$ by using **local Ward identities**
in a **dimensional regularization scheme** with $d = 3 - \varepsilon$.

After assuming the existence of a $O(1)$ fixed point
(suggested by a one loop calculation) they find that:

- * the condensate is stable in $2d$ and $T = 0$;
- * the theory displays a linear excitation spectrum a'la Bogoliubov.

- **Problems:**

- (1) Not even order by order results in $2d$ were available
- (2) The $2d$ theory is quite delicate: 8 effective couplings
(two of them relevant) and 1 free parameter

Two dimensions

- Pistolesi, Castellani, Di Castro, Strinati (1997 & 2004):
RG analysis in $3d$ and $2d$ by using **local Ward identities**
in a **dimensional regularization scheme** with $d = 3 - \varepsilon$.

After assuming the existence of a $O(1)$ fixed point
(suggested by a one loop calculation) they find that:

- * the condensate is stable in $2d$ and $T = 0$;
- * the theory displays a linear excitation spectrum a'la Bogoliubov.

► Problems:

- (1) Not even order by order results in $2d$ were available
- (2) The $2d$ theory is quite delicate: 8 effective couplings
(two of them relevant) and 1 free parameter
- (3) **the momentum cutoffs break the local gauge invariance**
 ↪ In low-dimensional systems of interacting fermions
 (**Luttinger liquids**) the corrections to WIs are crucial
 for establishing the infrared behavior of the system
 (Benfatto, Falco, Mastropietro, 2009)


The goal

Extend the Wilsonian RG approach to the Bose gas in the $2d$ continuum, at $T = 0$, both for $\rho_0 = 0$ and $\rho_0 > 0$, in the formalism developed by Benfatto and Gallavotti.


Exact RG approach

- Explicit bounds at all orders
- Complete control of all the diagrams (irrelevant terms included)
- Momentum cutoff regularization (essential for a perturbative construction)
- Corrections to Local Ward Identities can be studied within this scheme


Renormalization group results




Renormalization group results



Renormalization group results




Renormalization group results




¹Fisher, Weichmann, Grinstein, Fisher (1989), Sachdev, Senthil, Shankar (1994)


Renormalization group results



Renormalization group results



Renormalization group results



The functional integral representation

The interacting partition function can be formally expressed as a functional integral:

$$\frac{Z_\Lambda}{Z_\Lambda^0} = \int P_\Lambda^0(d\varphi) e^{-V_\Lambda(\varphi)}$$

- ▶ $\varphi_{\vec{x},t}^+ = (\varphi_{\vec{x},t}^-)^*$ complex fields (coherent states)
- ▶ $\Lambda = [0, \beta) \times L^2$

The functional integral representation

The interacting partition function can be formally expressed as a functional integral:

$$\frac{Z_\Lambda}{Z_\Lambda^0} = \int P_\Lambda^0(d\varphi) e^{-V_\Lambda(\varphi)}$$

- ▶ $\varphi_{\vec{x},t}^+ = (\varphi_{\vec{x},t}^-)^*$ complex fields (coherent states)
- ▶ $\Lambda = [0, \beta) \times L^2$

$$V_\Lambda(\varphi) = \frac{\lambda}{2} \int_{L^4} d^2\vec{x} d^2\vec{y} \int_{-\beta/2}^{\beta/2} dt |\varphi_{\vec{x},t}|^2 v(\vec{x}-\vec{y}) |\varphi_{\vec{y},t}|^2 - \bar{\nu}_{\beta,L} \int_{L^2} d^2\vec{x} \int_{-\beta/2}^{\beta/2} dt |\varphi_{\vec{x},t}|^2$$

The functional integral representation

The interacting partition function can be formally expressed as a functional integral:

$$\frac{Z_\Lambda}{Z_\Lambda^0} = \int P_\Lambda^0(d\varphi) e^{-V_\Lambda(\varphi)}$$

- ▶ $\varphi_{\vec{x},t}^+ = (\varphi_{\vec{x},t}^-)^*$ complex fields (coherent states)
- ▶ $\Lambda = [0, \beta) \times L^2$

$$V_\Lambda(\varphi) = \frac{\lambda}{2} \int_{L^4} d^2\vec{x} d^2\vec{y} \int_{-\beta/2}^{\beta/2} dt |\varphi_{\vec{x},t}|^2 v(\vec{x}-\vec{y}) |\varphi_{\vec{y},t}|^2 - \bar{v}_{\beta,L} \int_{L^2} d^2\vec{x} \int_{-\beta/2}^{\beta/2} dt |\varphi_{\vec{x},t}|^2$$

$P_\Lambda^0(d\varphi)$ is a complex Gaussian measure with covariance

$$S_\Lambda^0(x, y) = \langle a_x^+ a_y \rangle \Big|_{\lambda=0} = \int P_\Lambda^0(d\varphi) \varphi_x^- \varphi_y^+$$

$$\xrightarrow{|\Omega|, \beta \rightarrow \infty} \rho_0 + \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} d^2\vec{k} dk_0 \frac{e^{-ik \cdot x}}{-ik_0 + \vec{k}^2}$$

RG scheme for the BEC phase

$$\varphi_x^\pm = \xi^\pm + \psi_x^\pm \quad \text{with } \xi^\pm = |\Lambda|^{-1} \int_\Lambda \varphi_x^\pm dx, \quad \langle \xi^- \xi^+ \rangle = \rho_0, \quad \langle \psi_x^- \psi_x^+ \rangle \text{ decaying}$$

$$\frac{Z_\Lambda}{Z_\Lambda^0} = \int P_\Lambda^0(d\varphi) e^{-V_\Lambda(\varphi)}$$

RG scheme for the BEC phase

$$\varphi_x^\pm = \xi^\pm + \psi_x^\pm \quad \text{with } \xi^\pm = |\Lambda|^{-1} \int_\Lambda \varphi_x^\pm dx, \quad \langle \xi^- \xi^+ \rangle = \rho_0, \quad \langle \psi_x^- \psi_x^+ \rangle \text{ decaying}$$

$$\frac{Z_\Lambda}{Z_\Lambda^0} = \int P_\Lambda(d\xi) e^{-|\Lambda| f_\Lambda^B(\xi)} \int P_\Lambda(d\psi) e^{-Q_\Lambda(\psi, \xi) - \mathcal{V}_\Lambda^B(\psi, \xi)}$$

RG scheme for the BEC phase


$$\varphi_x^\pm = \xi^\pm + \psi_x^\pm \quad \text{with } \xi^\pm = |\Lambda|^{-1} \int_\Lambda \varphi_x^\pm dx, \quad \langle \xi^- \xi^+ \rangle = \rho_0, \quad \langle \psi_x^- \psi_x^+ \rangle \text{ decaying}$$

$$\frac{Z_\Lambda}{Z_\Lambda^0} = \int P_\Lambda(d\xi) e^{-|\Lambda| f_\Lambda^B(\xi)} \int P_\Lambda^B(d\psi) e^{-V_\Lambda^B(\psi, \xi)}$$

RG scheme for the BEC phase

$$\varphi_x^\pm = \xi^\pm + \psi_x^\pm \quad \text{with } \xi^\pm = |\Lambda|^{-1} \int_\Lambda \varphi_x^\pm dx, \quad \langle \xi^- \xi^+ \rangle = \rho_0,$$


$$\frac{Z_\Lambda}{Z_\Lambda^0} = \int P_\Lambda(d\xi) \underbrace{e^{-|\Lambda|f_\Lambda^B(\xi)} \int P_\Lambda^B(d\psi) e^{-V_\Lambda^B(\psi, \xi)}}_{e^{-|\Lambda|f_\Lambda(\xi)}}$$



RG scheme for the BEC phase

$$\varphi_x^\pm = \sqrt{\rho_0} + \psi_x^\pm$$






$$f(\rho) = f^B(\sqrt{\rho_0}) - \bar{\nu}\rho_0 + \lim_{|\Lambda| \rightarrow \infty} \frac{1}{|\Lambda|} \int P_\Lambda^B(d\psi) e^{-\bar{\nu}_\Lambda(\psi)}$$



RG scheme for the BEC phase

$$\varphi_x^\pm = \sqrt{\rho_0} + \psi_x^\pm$$






$$\Xi_\Lambda = \int P_\Lambda^B(d\psi) e^{-\bar{V}_\Lambda(\psi)}$$



RG scheme for the BEC phase

$$\varphi_x^\pm = \sqrt{\rho_0} + \psi_x^\pm$$

$$\Xi_\Lambda = \int P_\Lambda^B(d\psi) e^{-\tilde{V}_\Lambda(\psi)}$$



Bogoliubov propagator:


$$g^B(\mathbf{x} - \mathbf{y}) = \begin{pmatrix} g_{-+}^B(\mathbf{x} - \mathbf{y}) & g_{--}^B(\mathbf{x} - \mathbf{y}) \\ g_{++}^B(\mathbf{x} - \mathbf{y}) & g_{+-}^B(\mathbf{x} - \mathbf{y}) \end{pmatrix}$$

$$= \int \frac{dk_0 d^2 \vec{k}}{(2\pi)^3} \frac{e^{-i\vec{k}(\vec{x}-\vec{y}) - ik_0(x_0 - y_0)}}{k_0^2 + \varepsilon^2(\vec{k})} \begin{pmatrix} ik_0 + |\vec{k}|^2 + \lambda \hat{v}(\vec{k}) \rho_0 & -\lambda \rho_0 \hat{v}(\vec{k}) \\ -\lambda \rho_0 \hat{v}(\vec{k}) & -ik_0 + |\vec{k}|^2 + \lambda \hat{v}(\vec{k}) \rho_0 \end{pmatrix},$$

with $\varepsilon^2(\vec{k}) = |\vec{k}|^4 + 2\lambda \hat{v}(\vec{k}) \rho_0 |\vec{k}|^2$.


RG scheme for the BEC phase

$$\Xi_\Lambda = \int P_\Lambda^B(d\psi) e^{-\tilde{V}_\Lambda(\psi)}$$



RG scheme for the BEC phase


$$\Xi_\Lambda = e^{-|\Lambda|f_{\Lambda,h}} \int P_{\Lambda}^{\leq h}(d\psi) e^{-V_{\Lambda,h}(\psi)}$$



- 1 **Multiscale decomposition:** we integrate iteratively the fields of decreasing energy scale, e.g. $k_0^2 + 2^{\bar{h}} \vec{k}^2 \simeq 2^{2h}$ for $h \leq \bar{h}$.

RG scheme for the BEC phase

$$\Xi_\Lambda = e^{-|\Lambda|f_{\Lambda,h}} \int P_\Lambda^{\leq h}(d\psi) e^{-V_{\Lambda,h}(\psi)}$$




- 1 **Multiscale decomposition:** we integrate iteratively the fields of decreasing energy scale, e.g. $k_0^2 + 2^{\bar{h}} \vec{k}^2 \simeq 2^{2h}$ for $h \leq \bar{h}$.
- 2 Integration over the fields **higher than 2^h** : $V_h(\psi) = \mathcal{L}V_h(\psi) + \mathcal{R}V_h(\psi)$

$$\begin{aligned} \mathcal{L}V_h = & \text{[Diagram: Red crossed lines with } \lambda_h^6 \text{]} + \text{[Diagram: Black X with } 2^h \lambda_h \text{]} + \text{[Diagram: Black vertex with } 2^{\frac{h}{2}} \mu_h \text{]} + \text{[Diagram: Black line with } 2^{2h} \nu_h \text{]} \\ & + \text{[Diagram: Dashed line with } Z_h \text{]} + \text{[Diagram: Solid line with } \partial_{x_0} \text{ and } B_h \text{]} + \text{[Diagram: Solid line with } \partial_{x_i} \text{ and } A_h \text{]} + \text{[Diagram: Dashed line with } \partial_{x_0} \text{ and } E_h \text{]} \end{aligned}$$

RG scheme for the BEC phase

$$\Xi_\Lambda = e^{-|\Lambda|f_{\Lambda,h}} \int P_\Lambda^{\leq h}(d\psi) e^{-V_{\Lambda,h}(\psi)}$$




- 1 **Multiscale decomposition:** we integrate iteratively the fields of decreasing energy scale, e.g. $k_0^2 + 2^{\bar{h}} \vec{k}^2 \simeq 2^{2h}$ for $h \leq \bar{h}$.
- 2 Integration over the fields **higher than 2^h** : $V_h(\psi) = \mathcal{L}V_h(\psi) + \mathcal{R}V_h(\psi)$

$$\begin{aligned} \mathcal{L}V_h = & \text{diagram with } \lambda_h^6 \text{ and a red cross} + \text{diagram with } 2^h \lambda_h \text{ and a black dot} + \text{diagram with } 2^{\frac{h}{2}} \mu_h \text{ and a black dot} + \text{diagram with } 2^{2h} \nu_h \text{ and a black dot} \\ & + \text{diagram with } Z_h \text{ and a black dot} + \text{diagram with } B_h \text{ and a black dot} + \text{diagram with } A_h \text{ and a black dot} + \text{diagram with } E_h \text{ and a black dot} \end{aligned}$$

- 3 Using **the Gallavotti-Nicoló tree expansion** we prove that $\mathcal{R}V_h$ is well defined with explicit bounds if the terms in $\mathcal{L}V_h$ are bounded.

RG scheme for the BEC phase

$$\Xi_\Lambda = e^{-|\Lambda|f_{\Lambda,h}} \int P_\Lambda^{\leq h}(d\psi) e^{-V_{\Lambda,h}(\psi)}$$



- 1 **Multiscale decomposition:** we integrate iteratively the fields of decreasing energy scale, e.g. $k_0^2 + 2^{\bar{h}} \vec{k}^2 \simeq 2^{2h}$ for $h \leq \bar{h}$.
- 2 Integration over the fields **higher than 2^h** : $V_h(\psi) = \mathcal{L}V_h(\psi) + \mathcal{R}V_h(\psi)$

$$\begin{aligned} \mathcal{L}V_h = & \text{[Red vertex with } \lambda_h^6 \text{]} + \text{[Blue vertex with } 2^h \lambda_h \text{]} + \text{[Dashed blue vertex with } 2^{\frac{h}{2}} \mu_h \text{]} + \text{[Horizontal line with } 2^{2h} \nu_h \text{]} \\ & + \text{[Dashed blue vertex with } Z_h \text{]} + \text{[Horizontal line with } \frac{\partial_{x_0} \partial_{x_0}}{B_h} \text{]} + \text{[Horizontal line with } \frac{\partial_{x_i} \partial_{x_i}}{A_h} \text{]} + \text{[Horizontal line with } \frac{\partial_{x_0}}{E_h} \text{]} \end{aligned}$$

- 3 Using **the Gallavotti-Nicoló tree expansion** we prove that $\mathcal{R}V_h$ is well defined with explicit bounds if the terms in $\mathcal{L}V_h$ are bounded.
- 4 Cancellations in the beta function of Z_h and μ_h follow from **Global WIs**

Flow equations for the effective interactions *below* h^*


There are two effective (three and two body) interactions, whose flows are coupled among them at all orders. Under the assumptions on the propagator

$$E_h^2/(Z_h B_h) \ll 1$$


$$A_h/B_h = (\text{const.})$$

$$B_h \leq (\text{const.})$$

the flow equations for $x_h := \lambda_h$ and $y_h := \lambda_{6,h}/(\lambda_h^2)$ at leading order (in the continuum limit) are:



Numerical solutions to the leading order flows for x_h and y_h .



$$\begin{cases} \pi^2 \frac{dx}{dt} = \pi^2 x - 2x^2 \\ \pi^2 \frac{dy}{dt} = -2\pi^2 y + \frac{16}{3}x - 2xy + cx^2 \end{cases}$$

Flow equations for the effective interactions *below* h^*


There are two effective (three and two body) interactions, whose flows are coupled among them at all orders. Under the assumptions on the propagator

$$E_h^2/(Z_h B_h) \ll 1$$


$$A_h/B_h = (\text{const.})$$

$$B_h \leq (\text{const.})$$

the flow equations for $x_h := \lambda_h$ and $y_h := \lambda_{6,h}/(\lambda_h^2)$ at leading order (in the continuum limit) are:



Numerical solutions to the leading order flows for x_h and y_h .



$$\begin{cases} \pi^2 \frac{dx}{dt} &= \pi^2 x - 2x^2 \\ \pi^2 \frac{dy}{dt} &= -2\pi^2 y + \frac{16}{3}x - 2xy + cx^2 \end{cases}$$

Non perturbative fixed points?

Flow equations for the renormalized wave functions

Even assuming the existence of the fixed points, one is left with studying the flow of A_h , E_h and B_h .

- In Castellani et al. this was done by using **local Ward Identities**, checked with dimensional regularization at the one loop level.

Flow equations for the renormalized wave functions

Even assuming the existence of the fixed points, one is left with studying the flow of A_h , E_h and B_h .

- ▶ In Castellani et al. this was done by using **local Ward Identities**, checked with dimensional regularization at the one loop level.
- ▶ The arguments based on local WIs appears to be in disagreement with the one-loop computations within a momentum-cutoff scheme **also in the region where the theory is still perturbative.**

Flow equations for the renormalized wave functions

Even assuming the existence of the fixed points, one is left with studying the flow of A_h , E_h and B_h .

- ▶ In Castellani et al. this was done by using **local Ward Identities**, checked with dimensional regularization at the one loop level.
- ▶ The arguments based on local WIs appears to be in disagreement with the one-loop computations within a momentum-cutoff scheme **also in the region where the theory is still perturbative.**
- ! The source of the violation to the local WIs is a correction term due to the momentum cutoff.


Flow equations for the renormalized wave functions

Even assuming the existence of the fixed points, one is left with studying the flow of A_h , E_h and B_h .

- ▶ In Castellani et al. this was done by using **local Ward Identities**, checked with dimensional regularization at the one loop level.
- ▶ The arguments based on local WIs appears to be in disagreement with the one-loop computations within a momentum-cutoff scheme **also in the region where the theory is still perturbative.**
- ! **The source of the violation to the local WIs is a correction term due to the momentum cutoff.**

The findings based on the systematic use of local WIs, and then the nature and existence of the $2d$ condensate, should be seriously reconsidered.

Renormalization group results



Outlook

- ▶ Do corrections to LWIs correspond to anomalies ?

Outlook

- ▶ Do corrections to LWIs correspond to anomalies ?
- ▶ Different parameters regime in $2d$?


Outlook

- ▶ Do corrections to LWIs correspond to anomalies ?
- ▶ Different parameters regime in $2d$?
- ▶ Critical temperature
- ▶ ...
- ▶ Constructive theory (starting from the quantum critical point)

The superfluid-insulator transition in the boson Hubbard model¹

On site interacting bosons hoppings between sites i of a lattice,
 $t > 0$ (hopping parameter), $\lambda > 0$ and $\mu \leq 0$ (chemical potential):

$$H_L = -t \sum_{\langle ij \rangle} (\hat{a}_i^\dagger \hat{a}_j + \hat{a}_j^\dagger \hat{a}_i) - \mu \sum_i \hat{n}_i + \frac{\lambda}{2} \sum_i \hat{n}_i (\hat{n}_i - 1)$$



The chemical potential drives a Quantum Phase transition between:


- **Mott Insulator phase** with integer boson density (no phase coherence);
- **Superfluid phase** with delocalized wave function (long-range order).

¹Fisher, Weichmann, Grinstein, Fisher. Phys. Rev. **B 40**, 546 (1989)

The superfluid-insulator transition in the boson Hubbard model¹

On site interacting bosons hoppings between sites i of a lattice,
 $t > 0$ (hopping parameter), $\lambda > 0$ and $\mu \leq 0$ (chemical potential):

$$H_L = -t \sum_{\langle ij \rangle} (\hat{a}_i^\dagger \hat{a}_j + \hat{a}_j^\dagger \hat{a}_i) - \mu \sum_i \hat{n}_i + \frac{\lambda}{2} \sum_i \hat{n}_i (\hat{n}_i - 1)$$




The chemical potential drives a Quantum Phase transition between:

- **Mott Insulator phase** with integer boson density (no phase coherence);
- **Superfluid phase** with delocalized wave function (long-range order).

¹Fisher, Weichmann, Grinstein, Fisher. Phys. Rev. **B 40**, 546 (1989)

Gallavotti–Nicolò tree expansion

The $|h|$ -th step of the iterative integration can be graphically represented as a sum of trees over $|h|$ scale labels. The number n of endpoints represents the order in perturbation theory.



Gallavotti–Nicolò trees are a synthetic and convenient way to isolate the divergent terms, avoiding the problem of overlapping divergences.