



# Inverting an Imperfect Model



Tony O'Hagan

# Outline

---

- ▶ **The objectives of inversion**
  - ▶ Physical and tuning parameters
  - ▶ Imperfect simulators – two examples
- ▶ **Ignoring model discrepancy**
  - ▶ Results of inverting the examples
  - ▶ Implications for learning about physical parameters
- ▶ **The simple machine and model discrepancy**
  - ▶ Simple model discrepancy
  - ▶ Modelling discrepancy
  - ▶ Nonidentifiability
- ▶ **Conclusions**

# The Objectives of Inversion

# Inversion as nonlinear regression

---

- ▶ We have a simulator  $\eta(x, \theta)$  and observations

$$z_i = \eta(x_i, \theta) + \varepsilon_i$$

- ▶ In statistical language this is a nonlinear regression model
  - ▶ The inversion problem is one of inference about  $\theta$
- ▶ I'll be assuming the Bayesian paradigm
  - ▶ Requires a prior distribution for  $\theta$ 
    - ▶ Often assumed to be non-informative
  - ▶ Produces a posterior distribution
- ▶ Very common approach, but has a major flaw
  - ▶ The observations are of the real physical system  $\zeta(\cdot)$
  - ▶ And the simulator is invariably imperfect:  $\eta(\cdot, \theta) \neq \zeta(\cdot) \quad \forall \theta$

# Model discrepancy

---

▶ We should write

$$z_i = \zeta(x_i) + \varepsilon_i = \eta(x_i, \theta) + \delta(x_i) + \varepsilon_i$$

- ▶ where  $\delta(\cdot)$  is model discrepancy
- ▶ and is an unknown function

▶ Inference about  $\theta$  is now clearly more complex

- ▶ No longer just a nonlinear regression problem
- ▶ Some literature on correlated errors

▶ How important is it?

- ▶ That depends on the objectives of the inversion
- ▶ And in particular on the nature of  $\theta$

# Inversion and the nature of parameters


---

- ▶ Parameters may be physical or just for tuning
  - ▶ We adjust tuning parameters so the model fits reality better
    - ▶ We are not really interested in their 'true' values
  - ▶ Physical parameters are different
    - ▶ We are often really interested in true physical values
- ▶ What are we inverting for?
  - ▶ To learn about physical parameter values
    - ▶ Model discrepancy is hugely important and needs care and thought
  - ▶ To predict reality – within context and range of observations
    - ▶ Interpolation: model discrepancy is important but easily addressed
  - ▶ To predict reality – outside context/range of observations
    - ▶ Extrapolation: discrepancy hugely important, needs care and thought

# Example 1: A simple machine (SM)

- ▶ A machine produces an amount of work  $y$  which depends on the amount of effort  $x$  put into it
  - ▶ Ideally,  $y = \beta x$ 
    - ▶ Parameter  $\beta$  is the rate at which effort can be converted to work
    - ▶ It's a physical parameter
  - ▶ True value of  $\beta$  is  $\beta^* = 0.65$
- ▶ Graph shows observed data
  - ▶ Points lie below  $y = 0.65x$ 
    - ▶ For large enough  $x$
  - ▶ Because of losses due to friction etc.


$$\zeta(x) = 0.65x(1 + 0.05x)^{-1}$$



# Example 2: Hot and cold (HC)

---

- ▶ An object is placed in a hot medium
  - ▶ Initially it heats up but then cools as the medium cools
- ▶ Simulator
$$\eta(t, \theta) = \theta_1 t \exp(-\theta_2 t)$$
- ▶ Reality
$$\zeta(t) = 10t(1 + t^2/10)^{-1.5}$$
- ▶ Parameters are physical
  - ▶  $\theta_1$  is initial heating rate, a property of the object
  - ▶  $\theta_2$  controls the cooling, a property of the medium and setup
- ▶ Interested in parameters but also in
  - ▶ Maximum temperature  $\zeta_{\max}$  and time  $t_{\max}$  when max is reached





# Meaning of parameters

---

- ▶ What is the relationship between parameters and reality?
  - ▶ They don't appear in  $\zeta(\cdot)$
  - ▶ In the SM example,  $\beta$  is the gradient at the origin
    - ▶ Theoretical efficiency only achievable at low inputs
    - ▶ This is well defined for reality,  $\beta = 0.65$
  - ▶ In the HC example,  $\theta_1$  is the gradient at the origin
    - ▶ Again well defined,  $\theta_1 = 10$
  - ▶  $\theta_2$  is more difficult because in reality cooling is not exponential
    - ▶ We define  $\theta_2 = 0.413$  from log gradient at point of inflection
  - ▶  $\zeta_{\max}$  and  $t_{\max}$  are not really physical
    - ▶ From the simulator,  $\zeta_{\max} = \theta_1 \theta_2^{-1} e^{-1}$ ,  $t_{\max} = \theta_2^{-1}$
    - ▶ In reality,  $\zeta_{\max} = 12.172$  and  $t_{\max} = 2.236$  depend on  $\theta$  and the setup

# Ignoring model discrepancy

# SM assuming no discrepancy

---

- ▶ Following the usual approach, inversion is a simple matter of linear regression through the origin

$$z_i = \beta x_i + \varepsilon_i$$

- ▶ Here are some results from various sample sizes spread uniformly over 3 ranges of  $x$  values

Range	[0.1,2]	[0.1,4]	[2,6]
n=11	0.549 (0.063)	0.562 (0.029)	0.533 (0.023)
n=31	0.656 (0.038)	0.570 (0.017)	0.529 (0.011)
n=91	0.611 (0.021)	0.571 (0.012)	0.528 (0.007)
n infinite	0.605 (0)	0.565 (0)	0.529 (0)

# HC assuming no discrepancy

---

- ▶ These results are from samples of 91 observations over three different ranges
  - ▶ Almost every single posterior distribution is concentrated far from the true value

Range	[0.1,1]	[0.2,2]	[0.4,4]	TRUE
$\theta_1$	10.57 (0.11)	11.11 (0.13)	12.77 (0.19)	10
$\theta_2$	0.159 (0.049)	0.237 (0.023)	0.401 (0.033)	0.413
$t_{\max}$	5.00 (0.47)	3.61 (0.13)	2.52 (0.04)	2.24
$\zeta_{\max}$	19.42 (1.55)	14.75 (0.38)	11.85 (0.08)	12.17

# The problem is completely general

---

- ▶ Inverting (calibrating, tuning, matching) a wrong model gives parameter estimates that are wrong
  - ▶ Not equal to their true physical values – biased
  - ▶ With more data we become more sure of these wrong values
- ▶ The SM and HC are trivial models, but the same conclusions apply to all models
  - ▶ All models are wrong
  - ▶ In more complex models it is just harder to see what is going wrong
  - ▶ Even with the SM, it takes a lot of data to see any curvature in reality
- ▶ What can we do about this?

# The Simple Machine and Model Discrepancy

# SM revisited


---

- ▶ Kennedy and O'Hagan (2001) introduced discrepancy  $\delta(\cdot)$ 
  - ▶ Modelled it as a zero-mean Gaussian process
  - ▶ They claimed it acknowledges additional uncertainty
  - ▶ And mitigates against over-fitting of  $\theta$
- ▶ So add this model discrepancy term to the linear model of the simple machine

$$z_i = \beta x_i + \delta(x_i) + \varepsilon_i$$

- ▶ With  $\delta(\cdot)$  modelled as a zero-mean GP
- ▶ Posterior distribution of  $\beta$  now behaves quite differently
- ▶ Results here from extensive study of SM in
  - ▶ Brynjarsdóttir, J. and O'Hagan, A. (2014). Learning about physical parameters: The importance of model discrepancy. *Inverse Problems*, **30**, 114007 (24pp), November 2014.

# SM – inversion, with discrepancy




- ▶ Posterior distribution much broader and doesn't get worse with more data
  - ▶ But still misses the true value



# Interpolation

- ▶ Main benefit of simple GP model discrepancy is prediction


- ▶ E.g. at  $x = 1.5$



- ▶ Prediction within the range of the data is possible
  - ▶ And gets better with more data

# But when it comes to extrapolation ...

- ▶ ... at  $x = 6$



- ▶ More data doesn't help because it's all in the range  $[0, 4]$
- ▶ Prediction OK here but gets worse for larger  $x$

# Extrapolation

---

- ▶ One reason for wish to learn about physical parameters
  - ▶ Should be better for extrapolation than just tuning
- ▶ Without model discrepancy
  - ▶ The parameter estimates will be biased
  - ▶ Extrapolation will also be biased
    - ▶ Because best fitting parameter values are different in different parts of the control variable space
  - ▶ With more data we become more sure of these wrong values
- ▶ With GP model discrepancy
  - ▶ Extrapolating far from the data does not work
    - ▶ No information about model discrepancy
    - ▶ Prediction just uses the (calibrated) simulator

# We haven't solved the problem

---

- ▶ With simple GP model discrepancy the posterior distribution for  $\theta$  is typically much wider
  - ▶ Increases the chance that we cover the true value
  - ▶ But is not very helpful
  - ▶ And increasing data does not improve the precision
- ▶ Similarly, extrapolation with model discrepancy gives wide prediction intervals
  - ▶ And may still not be wide enough
- ▶ What's going wrong here?

# Nonidentifiability

---

- ▶ Formulation with model discrepancy is not identifiable
- ▶ For any  $\theta$ , there is a  $\delta(x)$  to match reality perfectly
  - ▶ Reality is  $r(x) = f(x, \theta) + \delta(x)$
  - ▶ Given  $\theta$  and  $r(x)$ , model discrepancy is  $\delta(x) = r(x) - f(x, \theta)$
- ▶ Suppose we had an unlimited number of observations
  - ▶ We would learn reality's true function  $r(x)$  exactly
    - ▶ Within the range of the data
    - ▶ Interpolation works
  - ▶ But we would still not learn  $\theta$ 
    - ▶ It could in principle be anything
  - ▶ And we would still not be able to extrapolate reliably

# The joint posterior

---

- ▶ Inversion leads to a joint posterior distribution for  $\theta$  and  $\delta(\mathbf{x})$
- ▶ But nonidentifiability means there are many equally good fits  $(\theta, \delta(\mathbf{x}))$  to the data
  - ▶ Induces strong correlation between  $\theta$  and  $\delta(\mathbf{x})$
  - ▶ This may be compounded by the fact that simulators often have large numbers of parameters
    - ▶ (Near-)redundancy means that different  $\theta$  values produce (almost) identical predictions
    - ▶ Sometimes called equifinality
- ▶ Within this set, the prior distributions for  $\theta$  and  $\delta(\mathbf{x})$  count

# The importance of prior information

---

- ▶ The nonparametric GP term allows the model to fit and predict reality accurately given enough data
  - ▶ Within the range of the data
- ▶ But it **doesn't** mean physical parameters are correctly estimated
  - ▶ The separation between original model and discrepancy is unidentified
  - ▶ Estimates depend on prior information
  - ▶ Unless the real model discrepancy is just the kind expected a priori the physical parameter estimates will still be biased
- ▶ To learn about  $\theta$  in the presence of model discrepancy we need better prior information
  - ▶ And this is also crucial for extrapolation

# Better prior information

---

## ▶ For calibration

- ▶ Prior information about  $\theta$  and/or  $\delta(x)$
- ▶ We wish to calibrate because prior information about  $\theta$  is not strong enough
- ▶ So prior knowledge of model discrepancy is crucial
  - ▶ In the range of the data

## ▶ For extrapolation


- ▶ All this plus good prior knowledge of  $\delta(x)$  *outside* the range of the calibration data
  - ▶ That's seriously challenging!
- ▶ In the SM, a model for  $\delta(x)$  that says it is zero at  $x = 0$ , then increasingly negative, should do better



# Inference about the physical parameter


► We conditioned the GP

- $\delta(0) = 0$
- $\delta'(0) = 0$
- $\delta'(0.5) < 0$
- $\delta'(1.5) < 0$




# Prediction

## Constr. GP prior on MD



$x = 1.5$

## Constr. GP prior on MD



$x = 6$

# Conclusions

# Summary

---

- ▶ **Without model discrepancy**
  - ▶ Inference about physical parameters will be wrong
    - ▶ And will get worse with more data
  - ▶ The same is true of prediction
    - ▶ Both interpolation and extrapolation
- ▶ **With crude GP model discrepancy**
  - ▶ Interpolation inference is OK
    - ▶ And gets better with more data
  - ▶ But we still get physical parameters and extrapolation wrong
- ▶ **The better our prior knowledge about model discrepancy**
  - ▶ The more chance we have of getting physical parameters right
  - ▶ Also extrapolation
    - ▶ But then we need even better prior knowledge