



koi Scrubs,
ultra stylish, ultra comfy!

SHOP NOW

ADVERTISEMENT

follow us on
twitter



August 27, 2014

Search

439 people on this site
powered by chartbeat

Other search tools: [Drugs](#) | [Health](#) | [News](#)

[Submit News](#) | [Contact Us](#) | [About](#)



16-25 Railcard

Get out there with 1/3 off rail fares

ONLY £30

BUY NOW

People from low socio-economic positions in Britain still eat more salt

Published on August 27, 2014 at 1:52 AM · No Comments

Print PDF

 Tweet 1
 Share

People from low socio-economic positions in Britain still eat more salt than those from higher socio-economic positions, irrespective of where they live.

A paper published in the *BMJ Open* journal and led by Warwick Medical School suggests social inequalities in salt intake have hardly changed in the period from 2000-01 to 2011. This is despite a national average salt reduction over this time.

This paper is the first to monitor social inequalities following the national salt reduction programme.

The research was carried out by the World Health Organization Collaborating Centre for Nutrition, based at Warwick Medical School, University of Warwick.

Researchers looked at the geographical distribution of habitual dietary salt intake in Britain and its association with manual occupations and educational attainments, both indicators of socio-economic position and key determinants of health.

The researchers used the British National Diet and Nutrition Survey (2008-11), a national representative sample of 1,027 men and women aged 19-64 years living in Britain. Salt intake was assessed with a 4-day food diary, assessing salt consumption from food. Salt added by consumers at the table and during cooking (discretionary salt) was not measured.

There was a significant reduction in dietary salt consumption from 2000-1 to 2008-11 of 0.9g of salt per day, consistent with the total reduction in salt consumption of 1.4g per day reported nationally when also discretionary use of salt is taken into account.

Professor Francesco Cappuccio, senior

Trending Stories

- Ebola virus outbreak: Equatorial Guinea temporarily suspends visas, regional flights
- Ragweed pollen reported for first time in 2014 allergy reporting season
- Vajrayana meditation linked with Tibetan Buddhism can enhance cognitive performance
- 'Soft' bipolar spectrum disorders differ from depression
- Acetyl fentanyl is more potent and dangerous than heroin
- [Microbes influence human](#)



16-25 Railcard

Get out there with 1/3 off rail fares

ONLY £30

BUY NOW

Advertisement



eating behavior, dietary choices



author and Director of the WHO Collaborating Centre, said: "Whilst we are pleased to record an average national reduction in salt consumption coming from food of nearly a gram per day, we are disappointed to find out that the benefits of such a programme have not reached those most in need. These results are important as people of low socio-economic background are more likely to develop [high blood pressure](#) (hypertension) and to suffer disproportionately from [strokes](#), [heart attacks](#) and [renal failure](#)."

Related Stories

- [Diet reduces inflammatory markers in 'real world' Type 2 diabetes](#)
- [Blood test can predict recurrence of HPV-linked oral cancers](#)
- [Neo40 nitric oxide supplement helps promote normal blood pressure](#)

Professor Cappuccio continued: "The diet of disadvantaged socio-economic groups tends to be made up of low-quality, salt-dense, high-fat, high-calorie unhealthy cheap foods. We have seen a reduction in salt intake in Britain thanks to a policy, which included awareness campaigns, food reformulation and monitoring. However, clearly poorer households still have less healthy shopping baskets and the broad reformulation of foods high in salt has not reached them as much as we would have hoped."

"In our continued effort to reduce population salt intake towards a 6g per day target in Britain, it is crucial to understand the reasons for these social inequalities so as to correct this gap for an equitable and cost-effective delivery of cardiovascular prevention", Professor Cappuccio concluded.

Source:

[University of Warwick](#)

Be the first to rate this post

Posted in: [Medical Research News](#) | [Healthcare News](#)

Tags: [Blood](#), [Blood Pressure](#), [Cardiovascular Disease](#), [Coronary Heart Disease](#), [Diet](#), [Disability](#), [Heart Disease](#), [High Blood Pressure](#), [Hypertension](#), [Medical School](#), [Nutrition](#), [Renal Failure](#), [Stroke](#)

[Permalink](#) | [Comments \(0\)](#)

Orange Standard, it's the new standard in affordable scrubs. [SHOP NOW](#)

▶ Latest News



[Unregulated trash burning around the globe pumps more air pollution](#)



[Meridian Health joins "The Baton Pass" to spread message of hope](#)

[Viewpoints: Check out those nursing home stars; Rove a](#)

[← Prior News](#)

[Scientists find that lack of naturally occurring protein linked to early signs of dementia](#)

[Next News →](#)

[Report tallies funds committed to federal insurance exchange](#)

Read in | [English](#) | [Español](#) | [Français](#) | [Deutsch](#) | [Português](#) | [Italiano](#) | [日本語](#) | [한국어](#) | [简体中文](#) | [繁體](#)