


Recent Results From H.E.S.S. -and a look at the future!




Paula Chadwick, Dept. of Physics
University of Durham

The Plan

- Some background information
- Recent H.E.S.S. results
 - The Galactic Plane survey
 - The Galactic Ridge
 - Dark matter searches
 - Starburst Galaxies
 - The PKS2155-30 flare and quantum gravity
 - Multiwavelength observations of M87
- CTA – the Cherenkov Telescope Array


Electromagnetic Spectrum




Satellite-based:
511 keV to
around 50 GeV

Ground-based:
~20 GeV+

Imaging
Atmospheric
Cherenkov
Technique



Important features of the technique.....



Excellent source location




Cannot observe during full moon




IACTs are *pointing* instruments



Clouds are bad!



Very large effective area



Energy threshold (and collection area) increase with zenith angle.

VHE Experimental World

MILAGRO



STACEE



MAGIC



TIBET



MILAGRO

STACEE

MAGIC

TIBET
ARGO-YBJ

TACTIC

PACT

GRAPES

VERITAS

TACTIC

HESS

CANGAROO III



Rene Ong

OG 1


HESS



CANGAROO



High Energy Stereoscopic System – H.E.S.S.



M-PIK Heidelberg; Humboldt University, Berlin; University of Hamburg; Ruhr University, Bochum; Landessternwarte Heidelberg; Tübingen University; Erlangen-Nürnberg University

LLR Ecole Polytechnique; LPNHE; APC College de France; University of Grenoble; CESR Toulouse; CEA Saclay; Observatoire de Paris-Meudon; LPTA Montpellier; LAPP Annecy

Durham University; University of Leicester

Dublin Institute for Advanced Studies

Polish Academy of Sciences (Astronomical Center & Institute of Nuclear Physics); Jagiellonian University; Nicolaus Copernicus University

Charles University, Prague

Yerevan Physics Institute, Armenia


University of Namibia

North-Western University, South Africa

University of Adelaide, Australia

University of Innsbruck, Austria

University of Stockholm, Sweden




System Parameters

- ⊕ Energy Threshold 100 GeV
- ⊕ Energy Resolution 15%
- ⊕ Field of View $\sim 5^\circ$
- ⊕ Angular Resolution 0.05° - 0.1°
- ⊕ Pointing Accuracy ~ 10 arcsec
- ⊕ Signal Rate $\sim 55/\text{min}$ (Crab Like)
- ⊕ Sensitivity:
 - ⊕ 1 Crab in 30 sec
 - ⊕ 0.01 Crab in 50h


(All at zenith)






Source Types

- Plerion PWN
- XRB PSR
- HBL IBL FRI FSRQ LBL
- Shell
- Starburst
- DARK
- MQS Cat. Var. UNID
Other BIN WR



Source Types

- Plerion PWN
- XRB PSR
- HBL IBL FRI FSRQ LBL
- Shell
- Starburst
- DARK
- MQS Cat. Var. UNID
Other BIN WR



Source Types

- Plerion PWN
- XRB PSR
- HBL IBL FRI FSRQ LBL
- Shell
- Starburst
- DARK
- MQS Cat. Var. UNID
Other BIN WR

Sources by Type


Unidentified	26 (and falling)	HBL	23
PWN	17	IBL	3
Shell SNRs	13	LBL	2
Binaries	5	FRI	2
Clusters/WR	3	Starburst Galaxies	2
Diffuse	2	FSRQ	1
		Gal. Centre	1 (!)

Fortuitously, that comes to 100 – but it's subjective!

Science with VHE Gamma Rays





Origin of
cosmic rays





SNRs

Space-time
& relativity




Pulsars
and PWN

Dark matter



AGNs


GRBs




Cosmology

The H.E.S.S. Galactic Plane Survey

The Extended H.E.S.S. GPS
2005 - 2008




Acceptance-corrected Exposure




Extended H.E.S.S. GPS

- $-85^\circ < l < 60^\circ$
- $-3^\circ < b < 3^\circ$
- Scan mode: 400 h
- Detected **50+ Galactic sources** of VHE gamma-rays
- ICRC 2007, DPG 2008, Gamma08





GC molecular clouds
Tsuboi et al. 1999



10 kyrs

H.E.S.S. Observations of Diffuse Emission in GC Region




Aharonian et al., *Nature*, 439, 695 (2006)

Top-down: Annihilation of dark matter particles


$$\chi \chi \rightarrow \gamma\gamma, \gamma Z, \gamma h$$




Matter distribution
expected to have
characteristic
density profile:
 $\sim r^{-1}$ (NFW)
to $r^{-1.5}$ (Moore)
sharp spike
with long tail
and characteristic
energy spectrum



Galactic Centre




Radio




TeV H.E.S.S.

DM Annihilation – angular distribution



Angular distribution of H.E.S.S. result consistent with a point source, once diffuse BG eliminated (16% of total emission). Assume a Gaussian centred on best-fit position → lower limit to slope of distribution -1.2 (i.e. cuspy)


DM annihilation - spectrum



The Position of the Galactic Centre Source

Radio contours of
Sgr A East (VLA)

Previous H.E.S.S.
best-fit centroid



First H.E.S.S. result was compatible with Sgr A East, Sgr A* and PWN candidate G359.95-0.04. Using paraxial optical cameras on telescopes reduced pointing errors from 20 arcsec to 6 arcsec per axis. Sgr A East looks to be ruled out as source of emission.

New H.E.S.S. best-fit centroid

Sgr Dwarf Spheroidal Galaxy

A deep-field Hubble Space Telescope image of the Sagittarius Dwarf Spheroidal Galaxy. The galaxy is a dense, elongated cluster of stars, appearing as a bright, irregular streak of light against the dark background of space. The stars are concentrated in the center and spread out towards the edges, with some individual stars appearing as bright, multi-colored points of light. The overall shape is roughly elliptical but with a complex, irregular distribution of stars.

Has crossed Milky Way
at least 10 times
without being
disrupted.

Good candidate for
substantial amount of
DM – not much gas, so
low CR background
too.


Handily, also off the
Galactic Plane.


Signal is expected to
come from a region
 ~ 1.5 pc, much smaller
than the H.E.S.S. PSF.
Profile (NFW...) doesn't matter!

HST Image

H.E.S.S. Observations

June 2006, 11 hours. Upper limit $E > 250$ GeV: $3.6 \times 10^{-12} \text{ cm}^{-2}\text{s}^{-1}$. (95% c.l.)






For core model, a lower limit for the $B^{(1)}$ mass of 500 GeV can be derived.

100h observation would enable the exclusion of much more pMSSM parameter space and all KK space for the core model


Canis Major 'Overdensity'





From Strasbourg Observatory

No Signal!



Mass of system not well known, so this is assuming mass of 3×10^8 solar masses.


The Electron Spectrum

The ATIC experiment observed a peak in the electron spectrum between 300 and 800 GeV.

Coupled with PAMELA excess, this has led to much speculation – e.g. dark matter, contribution from a local pulsar etc.


Measuring electron spectrum with a VHE gamma-ray experiment is tough – electrons and gamma rays both produce pure electromagnetic showers.

Have to use off-GP data and extensive simulations to derive an ‘electron likeness’ parameter, ζ .





H.E.S.S. Measurements

Overall electron flux is compatible with ATIC within errors, but H.E.S.S. data exclude presence of a pronounced peak in the electron spectrum, though an energy shift could be possible, so it cannot be definitively ruled out. However, it's hard to reconcile with a KK dark matter scenario.



Starburst Galaxies – why bother?

Compulsory
picture!



Starburst galaxies = lots of star formation (in a small region) = lots of supernovae = lots of particle (proton) acceleration + lots of gas = lots of VHE gamma rays = confirmation of suspicions about galactic CRs (and maybe information about galaxy/star formation)

NGC 253



$D = 3.9 \pm 0.4$ Mpc


SN rate $\sim 10\times$ Milky Way in
starburst region

Mean density of gas in
starburst region almost 10^3
higher than MW

Radio, thermal X-rays show
hot, diffuse halo consistent
with galactic wind

Discovered by Caroline
Herschel in 1783

H.E.S.S. Detection of NGC 253



Flux ($E > 220$ GeV): $5.5 \pm 1.0_{\text{stat}} \pm 2.8_{\text{sys}} \times 10^{-13} \text{ cm}^{-2}\text{s}^{-1}$


$\sim 0.3\%$ Crab flux

119 hours of observation

No evidence for variability

CR density in starburst region \sim
2000x that near the Solar
System, and ~ 1400 times that
near the GC

Fermi LAT detections of NGC253 & M82




Flux ($E > 100$ MeV):
 $1.6 \pm 0.5_{\text{stat}} \pm 0.3_{\text{sys}} \times 10^{-8} \text{ cm}^{-2}\text{s}^{-1}$

Flux ($E > 100$ MeV):
 $0.6 \pm 0.4_{\text{stat}} \pm 0.4_{\text{sys}} \times 10^{-8} \text{ cm}^{-2}\text{s}^{-1}$

No evidence for variability in either object


Interpretation I



Gamma-ray luminosity best correlates with SN rate *and* the mass of gas in the galaxy – perhaps not surprising.

BUT distribution of CRs is unlikely to be uniform – e.g. the GeV emission in LMC mostly comes from 30 Doradus and does not trace star formation & total gas mass.

Interpretation II




Emission models depend on many different parameters – agreement looks better for M82 than for NGC 253. In M82, the smooth power law connection between GeV & TeV emission suggests the same process produces both. Relationship less clear for NGC 253.


NGC 253 and Cosmic Rays

- 220 GeV generating protons need energy ~ 1300 GeV
- Given
 - CR energy production in equilibrium with losses from nuclear collisions;
 - Measured gas density and SN rate;
 - Production spectrum $\propto E^{-2.1}$
- Then calculate gamma ray flux to be factor of 10^2 higher than observed; suggests CRs in NGC 253 more likely to escape than expected
- NGC 253 is not a perfect CR ‘calorimeter’ – ISM does not act as a perfect ‘beam dump’
- Nevertheless, conversion efficiency of protons to gamma rays is still $\sim 10\times$ higher than in the Milky Way
- Starburst nucleus should outshine the rest of the galaxy (consistent with H.E.S.S. point source)

Interpretation III



M82




NGC 253

Assume protons (pion decay) gamma rays dominate

In M82: exploit uncertainties in SN explosion rate & efficiency of CR generation.

In NGC 253: exploit uncertainties in distance (2.5 Mpc has been quoted), diffusion timescales & cutoffs in the proton injection spectrum.


Active Galactic Nuclei



The most common VHE-emitting AGN are the high-frequency peaked blazars – where we are looking almost directly down the jet.



PKS2155-304 in 2006



In late July 2006, this AGN went crazy, and produced a burst that made the object 20 times brighter than the Crab Nebula. The burst contained over 60,000 gamma rays!


Energy Dependence of c


Broadly speaking (models vary), quantum gravity predicts an energy-dependence of the speed of light of the form:

$$c' = c \left(1 + \xi \frac{E}{E_p} + \zeta \frac{E^2}{E_p^2} \right)$$

where E_p is the Planck Energy, 1.22×10^{19} GeV, and ξ and ζ are free parameters to be determined. The correction is expected to be very small, but Amelino-Camelia et al. (1998) suggested that these modifications can produce significant time delays with energy over cosmological distances. The absence of such energy dispersion sets limits on ξ and ζ .

We can use the massive flare from PKS2155-304 to test this.





The MCCF (left) looks quite exciting, with an apparent 20s lag for higher energy. However, when you do 10,000 simulations varying the flux points of the oversampled light curve within measurement errors and create a cross-correlation peak distribution (right), you find an RMS of 28s and that simulations produce a negative delay for 21% of the time. The 'lag' is therefore consistent with zero.

$$|\xi| < 17 \text{ for linear dispersion \& } |\zeta| < 7.3 \times 10^{19} \text{ for quadratic dispersion}$$

M87 – a Radio Galaxy




Pinpointing the Emission Site


Combined observations of H.E.S.S., VERITAS, MAGIC & the VLBA 43 GHz team – the paper has 392 authors!

Emission is variable on \sim day timescales. Previous observations had shown an increase in VHE emission roughly contemporaneous with emission from the knot region HST-1 in the jet.

These observations show the emission is coming from the nucleus.




Which came first??



The core seems to show a period of below-normal activity before the flare, and the radio flux increase actually starts before the VHE flare. This is followed by enhanced emission along the inner jet after the VHE flare.

adapted by Hinton from
Horan & Weekes 2003



- Current instruments have passed the critical sensitivity threshold and reveal a rich panorama, **but this is clearly only the tip of the iceberg**



So what next???

The Cherenkov Telescope Array (CTA) a 'real' observatory with ~ 100 telescopes

25 MEuro


Low-energy:
energy threshold
of few 10 GeV

35 MEuro

Core array:
mCrab sensitivity at
100 GeV–10 TeV

20 MEuro

High-energy:
10 km² area at
multi-TeV energies



Work Packages

WP1	MNG	Management of the design study
WP2	PHYS	Astrophysics and astroparticle physics
WP3	MC	Optimization of array layout, performance studies, ...
WP4	SITE	Site evaluation and site infrastructure
WP5	MIR	Telescope optics, mirrors, mirror alignment
WP6	TEL	Telescope structure, drive, control, robotics
WP7	FPI	Focal plane instrumentation, mechanics and photo detectors
WP8	ELEC	Readout electronics and trigger
WP9	ATAC	Atmospheric monitoring, associated science & instrument calib.
WP10	OBS	Observatory operation and access
WP11	DATA	Data handling, data processing, data management and access
WP12	QA	Risk assessment and quality assurance, production planning

UK well represented
↓

D Torres

J Hinton

G Vasileiadis

M Mariotti & M Doro

M Panter

R Mirzoyan

P Vincent

S Nolan

A Sillanpää & S Wagner

C Stegmann

M Punch & M Benallou

J Knapp

(acting) Chair of the Consortium Board

FP7 Preparatory Phase:

SST: small-size telescopes

MST: medium-size telescopes

LST: large-size telescope

UK emphasis ←

Opportunities for UK contributions,
Manpower needed everywhere.

Stolen from Johannes Knapp!

Funding:

"guaranteed"

ASPERA Common Call CTA

mostly personnel

€ 2.6M

UK: ~~£0.5M~~

£0.00

FP7 CTA Prep Phase call (EU) (€ 6M)
announcement spring 2010

UK: ≈ € 0.78M

mostly organisational

matching funds:

€ 2.93M

UK: ≈ € 0.26M

FP7 Virtual research infrastructures (EU) (€ 4.2M)

announcement spring 2010 UK: ≈ 4 PD yrs

GRID, archiving, data handling, ...

other funding :

€ 2.76M

UK: ≈ € 0.24M

